

## 3.2 Reflection and Refraction

- Geometrical Optics
- Reflection
- Refraction




Christian Huygens


## Geometrical optics

In geometrical optics light waves are considered to move in straight lines. This is a good description as long as the waves do not pass through small openings (compared to  $\lambda$ )

## Light waves



Rays are perpendicular to wave fronts




## Reflection

- Two general types of reflection
  - Specular reflection
  - Diffuse reflection
- Most of geometric optics deals with specular reflection.
- However, most of the time ambient lighting is due to diffuse reflection.


### Specular reflection

Flat surface  
Light reflected in one direction





### Diffuse reflection

Rough surface  
Light reflected in all directions



## Transmission and Reflection at an interface



What are some examples of these processes in this picture.

Specular Reflection

Diffuse reflection (scattering)


Transmission

Absorption

## Law of Reflection

The angle of reflection equals the angle of incidence


$$\theta_1 = \theta_1'$$



## Multiple reflections

- For multiple reflections use the law of reflection for each reflecting surface.

## 2-Dimensional Corner reflector



## Refraction

- Refraction is the bending of light when it passes across an interface between two materials.
- The bending is due to the differences in the speed of light in different media.
- The index of refraction of a material  $n_i$  is the ratio of the speed of light in a vacuum  $c$  to the speed of light in the material  $v_i$

$$n_i = \frac{c}{v_i}$$

**TABLE 22.1**  
Indices of Refraction for Various Substances, Measured with Light of Vacuum Wavelength  $\lambda_0 = 589 \text{ nm}$

Substance	Index of Refraction	Substance	Index of Refraction
<b>Solids at 20°C</b>		<b>Liquids at 20°C</b>	
Diamond (C)	2.419	Benzene	1.501
Fluorite ( $\text{CaF}_2$ )	1.434	Carbon disulfide	1.628
Fused quartz ( $\text{SiO}_2$ )	1.458	Carbon tetrachloride	1.461
Glass, crown	1.52	Ethyl alcohol	1.361
Glass, flint	1.66	Glycerine	1.473
Ice ( $\text{H}_2\text{O}$ ) (at 0°C)	1.309	Water	1.333
Polystyrene	1.49		
Sodium chloride ( $\text{NaCl}$ )	1.544	<b>Gases at 0°C, 1 atm</b>	
Zircon	1.923	Air	1.000 29
		Carbon dioxide	1.000 45

The speed of light changes because of absorption and emission of light by atoms in the media which produce a time lag.

### Transmission across an interface

The speed of the wave changes.  
The frequency remains the same.  
The wavelength changes

### Refraction and Reflection

### Snell's Law of Refraction

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

Going from air to glass

$n_2 > n_1$   
 $\theta_2 < \theta_1$

( $\sin \theta$  increases with  $\theta$ )

Going from glass to air


$n_2 < n_1$   
 $\theta_2 > \theta_1$

### Physical picture for Snell's Law

One end of the wave front slows down.  
The wave front changes direction.

### Example 22.2

Find the angle of refraction for an angle of incidence of  $30^\circ$  in going from air to glass ( $n_{\text{glass}} = 1.52$ )



### Example 22.4

Show that light going through a flat slab is not deviated in angle.

