


7-1.
$$E_{n_1 n_2 n_3} = \frac{\hbar^2 \pi^2}{2mL^2} (n_1^2 + n_2^2 + n_3^2) \quad (\text{Equation 7-4})$$

$$E_{311} = \frac{\hbar^2 \pi^2}{2mL^2} (3^2 + 1^2 + 1^2) = 11E_0 \quad \text{where } E_0 = \frac{\hbar^2 \pi^2}{2mL^2}$$

$$E_{222} = E_0 (2^2 + 2^2 + 2^2) = 12E_0 \quad \text{and} \quad E_{321} = E_0 (3^2 + 2^2 + 1^2) = 14E_0$$

The 1st, 2nd, 3rd, and 5th excited states are degenerate.



$$7-2. \quad E_{n_1 n_2 n_3} = \frac{\hbar^2 \pi^2}{2m} \left(\frac{n_1^2}{L_1^2} + \frac{n_2^2}{L_2^2} + \frac{n_3^2}{L_3^2} \right) = \frac{\hbar^2 \pi^2}{2mL_1^2} \left(n_1^2 + \frac{n_2^2}{4} + \frac{n_3^2}{9} \right) \quad (\text{Equation 7-5})$$

$n_1 = n_2 = n_3 = 1$ is the lowest energy level.

$$E_{111} = E_0 \left(1 + 1/4 + 1/9 \right) = 1.361E_0 \quad \text{where } E_0 = \frac{\hbar^2 \pi^2}{2mL_1^2}$$

The next nine levels are, increasing order,

(Problem 7-2 continued)

n_1	n_2	n_3	$E (\times E_0)$
1	1	2	1.694
1	2	1	2.111
1	1	3	2.250
1	2	2	2.444
1	2	3	3.000
1	1	4	3.028
1	3	1	3.360
1	3	2	3.472
1	2	4	3.778

7-3. (a) $\psi_{n_1 n_2 n_3}(x, y, z) = A \cos \frac{n_1 \pi x}{L} \sin \frac{n_2 \pi y}{L} \sin \frac{n_3 \pi z}{L}$

(b) They are identical. The location of the coordinate origin does not affect the energy

level structure.

7-4. $\psi_{111}(x, y, z) = A \sin \frac{\pi x}{L_1} \sin \frac{\pi y}{2L_1} \sin \frac{\pi z}{3L_1}$

$$\psi_{112}(x, y, z) = A \sin \frac{\pi x}{L_1} \sin \frac{\pi y}{2L_1} \sin \frac{2\pi z}{3L_1}$$

$$\psi_{121}(x, y, z) = A \sin \frac{\pi x}{L_1} \sin \frac{\pi y}{L_1} \sin \frac{\pi z}{3L_1}$$

$$\psi_{122}(x, y, z) = A \sin \frac{\pi x}{L_1} \sin \frac{\pi y}{L_1} \sin \frac{2\pi z}{3L_1}$$

$$\psi_{113}(x, y, z) = A \sin \frac{\pi x}{L_1} \sin \frac{\pi y}{2L_1} \sin \frac{\pi z}{L_1}$$

7-5.

$$E_{n_1 n_2 n_3} = \frac{\hbar^2 \pi^2}{2m} \left(\frac{n_1^2}{L_1^2} + \frac{n_2^2}{(2L_1)^2} + \frac{n_3^2}{(4L_1)^2} \right) = \frac{\hbar^2 \pi^2}{2mL_1^2} \left(n_1^2 + \frac{n_2^2}{4} + \frac{n_3^2}{16} \right) \quad (\text{from Equation 7-5})$$

$$E_0 = \left(n_1^2 + \frac{n_2^2}{4} + \frac{n_3^2}{16} \right) \quad \text{where } E_0 = \frac{\hbar^2 \pi^2}{2mL_1^2}$$

(Problem 7-5 continued)

(a)

n_1	n_2	n_3	$E (\times E_0)$
1	1	1	1.313
1	1	2	1.500
1	1	3	1.813
1	2	1	2.063
1	1	4	2.250
1	2	2	2.250
1	2	3	2.563
1	1	5	2.813
1	2	4	3.000
1	1	6	3.500

(b) 1,1,4 and 1,2,2

$$7-7. \quad E_0 = \frac{\hbar^2 \pi^2}{2mL^2} = \frac{(1.055 \times 10^{-34} \text{ J}\cdot\text{s})^2 \pi^2}{2(9.11 \times 10^{-31} \text{ kg})(0.10 \times 10^{-9} \text{ m})^2 (1.609 \times 10^{-19} \text{ J/eV})} = 37.68 \text{ eV}$$

$$E_{311} - E_{111} = \Delta E = 11E_0 - 3E_0 = 8E_0 = 301 \text{ eV}$$

(Problem 7-7 continued)

$$E_{222} - E_{111} = \Delta E = 12E_0 - 3E_0 = 9E_0 = 339 \text{ eV}$$

$$E_{321} - E_{111} = \Delta E = 14E_0 - 3E_0 = 11E_0 = 415eV$$

7-8. (a) Adapting Equation 7-3 to two dimensions (i.e., setting $k_3 = 0$), we have

$$\psi_{n_1 n_2} = A \sin \frac{n_1 \pi x}{L} \sin \frac{n_2 \pi y}{L}$$

(b) From Equation 7-5, $E_{n_1 n_2} = \frac{\hbar^2 \pi^2}{2mL^2} (n_1^2 + n_2^2)$

(c) The lowest energy degenerate states have quantum numbers $n_1 = 1, n_2 = 2$,
and $n_1 = 2,$

$$n_2 = 1.$$

7-9. (a) For $n = 3$, $\ell = 0, 1, 2$

(b) For $\ell = 0, m = 0$. For $\ell = 1, m = -1, 0, +1$. For $\ell = 2, m = -2, -1, 0, +1, +2$.

(c) There are nine different m -states, each with two spin states, for a total of 18 states for

$$n = 3.$$

7-10. (a) For $\ell = 4$

$$L = \sqrt{\ell(\ell+1)}\hbar = \sqrt{4(5)}\hbar = \sqrt{20}\hbar$$

$$m_\ell = 4\hbar$$

$$\theta_{\min} = \cos^{-1} \frac{4}{\sqrt{20}} \rightarrow \theta_{\min} = 26.6^\circ$$


(b) For $\ell = 2$

$$L = \sqrt{6}\hbar \quad m_\ell = 2\hbar$$


$$\theta_{\min} = \cos^{-1} \frac{2}{\sqrt{6}} \rightarrow \theta_{\min} = 35.3^\circ$$

7-12. (a)

$$\ell = 1$$
$$|\mathbf{L}| = \sqrt{2}\hbar$$



(b)




$$\ell = 2$$

$$|\mathbf{L}| = \sqrt{6}\hbar$$

(c)

$$\ell = 4$$

$$|\mathbf{L}| = \sqrt{20}\hbar$$



(d) $|\mathbf{L}| = \sqrt{\ell(\ell+1)}\hbar$ (See diagrams above.)

$$7-13. \quad L^2 = L_x^2 + L_y^2 + L_z^2 \rightarrow L_x^2 + L_y^2 = L^2 - L_z^2 = \ell(\ell+1)\hbar^2 - (m\hbar)^2 = (6 - m^2)\hbar^2$$

$$(a) \quad (L_x^2 + L_y^2)_{\min} = (6 - 2^2)\hbar^2 = 2\hbar^2$$

$$(b) \quad (L_x^2 + L_y^2)_{\max} = (6 - 0^2)\hbar^2 = 6\hbar^2$$

(c) $L_x^2 + L_y^2 = (6-1)\hbar^2 = 5\hbar^2$ L_x and L_y cannot be determined separately.

(d) $n = 3$

7-15. $\mathbf{L} = \mathbf{r} \times \mathbf{p}$ $\frac{d\mathbf{L}}{dt} = \frac{d\mathbf{r}}{dt} \times \mathbf{p} + \mathbf{r} \times \frac{d\mathbf{p}}{dt}$

$\frac{d\mathbf{r}}{dt} \times \mathbf{p} = \mathbf{v} \times m\mathbf{v} = m\mathbf{v} \times \mathbf{v} = 0$ and $\mathbf{r} \times \frac{d\mathbf{p}}{dt} = \mathbf{r} \times \mathbf{F}$. Since for $V = V(r)$, i.e., central forces,

\mathbf{F} is parallel to \mathbf{r} , then $\mathbf{r} \times \mathbf{F} = 0$ and $\frac{d\mathbf{L}}{dt} = 0$

7-16. (a) For $\ell = 3$, $n = 4, 5, 6, \dots$ and $m = -3, -2, -1, 0, 1, 2, 3$

(b) For $\ell = 4$, $n = 5, 6, 7, \dots$ and $m = -4, -3, -2, -1, 0, 1, 2, 3, 4$

(c) For $\ell = 0$, $n = 1$ and $m = 0$

(d) The energy depends only on n . The minimum in each case is:

$$E_4 = -13.6eV / n^2 = -13.6eV / 4^2 = -0.85eV$$

$$E_5 = -13.6eV / 5^2 = -0.54eV$$

$$E_1 = -13.6eV$$

7-17. (a) $6f$ state: $n = 6, \ell = 3$

(b) $E_6 = -13.6eV / n^2 = -13.6eV / 6^2 = -0.38eV$

(c) $L = \sqrt{\ell(\ell+1)}\hbar = \sqrt{3(3+1)}\hbar = \sqrt{12}\hbar = 3.65 \times 10^{-34} \text{ J}\cdot\text{s}$

(d) $L_z = m\hbar$ $L_z = -3\hbar, -2\hbar, -1\hbar, 0, 1\hbar, 2\hbar, 3\hbar$

7-20. (a) For the ground state, $P(r)\Delta r = \psi^2 (4\pi r^2) \Delta r = \frac{4r^2}{a_0^3} e^{-2r/a_0} \Delta r$

For $\Delta r = 0.03a_0$, at $r = a_0$ we have $P(r)\Delta r = \frac{4a_0^2}{a_0^3} e^{-2} (0.03a_0) = 0.0162$

(b) For

$\Delta r = 0.03a_0$, at $r = 2a_0$ we have $P(r)\Delta r = \frac{4(2a_0)^2}{a_0^3} e^{-4} (0.03a_0) = 0.0088$

7-21. $P(r) = Cr^2 e^{-2Zr/a_0}$ For $P(r)$ to be a maximum,

$$\frac{dP}{dr} = C \left[r^2 \left(-\frac{2Z}{a_0} \right) e^{-2Zr/a_0} + 2r e^{-2Zr/a_0} \right] = 0 \rightarrow C \times \frac{2Zr}{a_0} \left(\frac{a_0}{Z} - r \right) e^{-2Zr/a_0} = 0$$

This condition is satisfied with $r = 0$ or $r = a_0/Z$. For $r = 0$, $P(r) = 0$ so the maximum

$P(r)$ occurs for $r = a_0/Z$.

7-22.
$$\int_0^\infty \int_0^\pi \int_0^{2\pi} \psi^2 r^2 \sin\theta dr d\theta d\phi = 1$$

$$= 4\pi \int_0^\infty \psi^2 r^2 dr = 4\pi C_{210}^2 \int_0^\infty \left(\frac{Zr}{a_0} \right)^2 r^2 e^{-Zr/a_0} dr = 1$$

$$= 4\pi C_{210}^2 \int_0^\infty \left(\frac{Z^2 r^4}{a_0^2} \right) e^{-Zr/a_0} dr = 1$$

Letting $x = Zr/a_0$, we have that $r = a_0 x/Z$ and $dr = a_0 dx/Z$ and

substituting

these above,

$$\int \psi^2 d\tau = \frac{4\pi a_0^3 C_{210}^2}{Z^3} \int_0^\infty x^4 e^{-x} dx$$

Integrating on the right side

$$\int_0^\infty x^4 e^{-x} dx = 6$$

Solving for C_{210}^2 yields:
$$C_{210}^2 = \frac{Z^3}{24\pi a_0^3} \rightarrow C_{210} = \left(\frac{Z^3}{24\pi a_0^3} \right)^{1/2}$$

7-26. For the most likely value of r , $P(r)$ is a maximum, which requires that (see Problem 7-24)

$$\frac{dP}{dr} = A \cos^2 \theta \left[r^4 \left(-\frac{Z}{a_0} \right) e^{-Zr/a_0} + 4r^3 e^{-Zr/a_0} \right] = 0$$

For hydrogen $Z = 1$ and $A \cos^2 \theta \left(r^3 / a_0 \right) (4a_0 - r) e^{-r/a_0} = 0$. This is satisfied for $r = 0$

and $r = 4a_0$. For $r = 0$, $P(r) = 0$ so the maximum $P(r)$ occurs for $r = 4a_0$.

7-33. (a) There should be four lines corresponding to the four m_j values $-3/2, -1/2, +1/2, +3/2$.

(b) There should be three lines corresponding to the three m_ℓ values $-1, 0, +1$.

7-68.
$$P(r) = \frac{4Z^3}{a_0^3} r^2 e^{-2Zr/a_0} \quad (\text{See Problem 7-63})$$

For hydrogen, $Z = 1$ and at the edge of the proton $r = R_0 = 10^{-15} m$. At that point, the

exponential factor in $P(r)$ has decreased to:

$$e^{-2R_0/a_0} = e^{-2(10^{-15})/(0.529 \times 10^{-10} m)} = e^{-(3.78 \times 10^{-5})} \approx 1 - 3.78 \times 10^{-5} \approx 1$$

Thus, the probability of the electron in the hydrogen ground state being inside the nucleus,

to better than four figures, is:

$$\begin{aligned} P(r) &= \frac{4r^2}{a_0^3} & P &= \int_0^{R_0} P(r) dr = \int_0^{R_0} \frac{4r^2}{a_0^3} = \frac{4}{a_0^3} \int_0^{R_0} r^2 dr = \frac{4}{a_0^3} \frac{r^3}{3} \Big|_0^{R_0} \\ & & &= \frac{4}{a_0^3} \left(\frac{R_0^3}{3} \right) = \frac{4(10^{-15} m)^3}{3(0.529 \times 10^{-10} m)^3} = 9.0 \times 10^{-15} \end{aligned}$$

7-70. (a) Substituting $\psi(r, \theta)$ into Equation 7-9 and carrying out the indicated operations

yields (eventually):

$$-\frac{\hbar^2}{2\mu} \psi(r, \theta) \left[2/r^2 - 1/4a_0^2 \right] - \frac{\hbar^2}{2\mu} \psi(r, \theta) (-2/r^2) + V\psi(r, \theta) = E\psi(r, \theta)$$

Canceling $\psi(r, \theta)$ and recalling that $r^2 = 4a_0^2$ (because ψ given is for $n = 2$)

we

$$\text{have } -\frac{\hbar^2}{2\mu}(-1/4a_0^2) + v = E$$

The circumference of the $n = 2$ orbit is:

$$C = 2\pi(4a_0) = 2\lambda \rightarrow a_0 = \lambda/4\pi = 1/2k.$$

$$\text{Thus, } -\frac{\hbar^2}{2\mu}\left(-\frac{1}{4/4k^2}\right) + V = E \rightarrow \frac{\hbar^2 k^2}{2\mu} + V = E$$

(b) or $\frac{p^2}{2m} + v = E$ and Equation 7-9 is satisfied.

$$\int_0^\infty \psi^2 dx = \int A^2 \left(\frac{r}{a_0}\right)^2 e^{-r/a_0} \cos^2 \theta r^2 \sin \theta dr d\theta d\phi = 1$$

$$A^2 \int_0^\infty \left(\frac{r}{a_0}\right)^2 e^{-r/a_0} r^2 dr \int_0^\pi \cos^2 \theta \sin \theta d\theta \int_0^{2\pi} d\phi = 1$$

Integrating (see Problem 7-22),

$$A^2 (6a_0^3)(2/3)(2\pi) = 1$$

$$A^2 = 1/8a_0^3\pi \rightarrow A = \sqrt{1/8a_0^3\pi}$$