

6 Interacting Classical Systems : Worked Examples

(6.1) Consider a model in which there are three possible states per site, which we can denote by A, B, and V. The states A and B are for our purposes identical. The energies of A-A, A-B, and B-B links are all identical and equal to W . The state V represents a vacancy, and any link containing a vacancy, meaning A-V, B-V, or V-V, has energy 0.

- (a) Suppose we write $\sigma = +1$ for A, $\sigma = -1$ for B, and $\sigma = 0$ for V. How would you write a Hamiltonian for this system? Your result should be of the form

$$\hat{H} = \sum_{\langle ij \rangle} E(\sigma_i, \sigma_j) \quad .$$

Find a simple and explicit function $E(\sigma, \sigma')$ which yields the correct energy for each possible bond configuration.

- (b) Consider a triangle of three sites. Find the average total energy at temperature T . *There are $3^3 = 27$ states for the triangle. You can just enumerate them all and find the energies.*
- (c) For a one-dimensional ring of N sites, find the 3×3 transfer matrix R . Find the free energy per site $F(T, N)/N$ and the ground state entropy per site $S(T, N)/N$ in the $N \rightarrow \infty$ limit for the cases $W < 0$ and $W > 0$. Interpret your results. *The eigenvalue equation for R factorizes, so you only have to solve a quadratic equation.*

Solution :

- (a) The quantity σ_i^2 is 1 if site i is in state A or B and is 0 in state V. Therefore we have

$$\hat{H} = W \sum_{\langle ij \rangle} \sigma_i^2 \sigma_j^2 \quad .$$

(b) Of the 27 states, eight have zero vacancies – each site has two possible states A and B – with energy $E = 3W$. There are 12 states with one vacancy, since there are three possible locations for the vacancy and then four possibilities for the remaining two sites (each can be either A or B). Each of these 12 single vacancy states has energy $E = W$. There are 6 states with two vacancies and 1 state with three vacancies, all of which have energy $E = 0$. The partition function is therefore

$$Z = 8 e^{-3\beta W} + 12 e^{-\beta W} + 7 \quad .$$

Note that $Z(\beta = 0) = \text{Tr } 1 = 27$ is the total number of ‘microstates’. The average energy is then

$$E = -\frac{1}{Z} \frac{\partial Z}{\partial \beta} = \left(\frac{24 e^{-3\beta W} + 12 e^{-\beta W}}{8 e^{-3\beta W} + 12 e^{-\beta W} + 7} \right) W$$

- (c) The transfer matrix is

$$R_{\sigma\sigma'} = e^{-\beta W \sigma^2 \sigma'^2} = \begin{pmatrix} e^{-\beta W} & e^{-\beta W} & 1 \\ e^{-\beta W} & e^{-\beta W} & 1 \\ 1 & 1 & 1 \end{pmatrix} \quad ,$$

where the row and column indices are A (1), B (2), and V (3), respectively. The partition function on a ring of N sites is

$$Z = \lambda_1^N + \lambda_2^N + \lambda_3^N \quad ,$$

where $\lambda_{1,2,3}$ are the three eigenvalues of R . Generally the eigenvalue equation for a 3×3 matrix is cubic, but we can see immediately that $\det R = 0$ because the first two rows are identical. Thus, $\lambda = 0$ is a solution to the characteristic equation $P(\lambda) = \det(\lambda \mathbb{I} - R) = 0$, and the cubic polynomial $P(\lambda)$ factors into the product of λ and a quadratic. The latter is easily solved. One finds

$$P(\lambda) = \lambda^3 - (2x + 1)\lambda^2 + (2x - 2)\lambda \quad ,$$

where $x = e^{-\beta W}$. The roots are $\lambda = 0$ and

$$\lambda_{\pm} = x + \frac{1}{2} \pm \sqrt{x^2 - x + \frac{9}{4}} \quad .$$

The largest of the three eigenvalues is λ_+ , hence, in the thermodynamic limit,

$$F = -k_B T \ln Z = -N k_B T \ln \left(e^{-W/k_B T} + \frac{1}{2} + \sqrt{e^{-2W/k_B T} - e^{-W/k_B T} + \frac{9}{4}} \right) \quad .$$

The entropy is $S = -\frac{\partial F}{\partial T}$. In the limit $T \rightarrow 0$ with $W < 0$, we have

$$\lambda_+(T \rightarrow 0, W < 0) = 2 e^{|W|/k_B T} + e^{-|W|/k_B T} + \mathcal{O}(e^{-2|W|/k_B T}) \quad .$$

Thus

$$\begin{aligned} F(T \rightarrow 0, W < 0) &= -N |W| - N k_B T \ln 2 + \dots \\ S(T \rightarrow 0, W < 0) &= N \ln 2 \quad . \end{aligned}$$

When $W > 0$, we have

$$\lambda_+(T \rightarrow 0, W > 0) = 2 + \frac{2}{3} e^{-W/k_B T} + \mathcal{O}(e^{-2W/k_B T}) \quad .$$

Then

$$\begin{aligned} F(T \rightarrow 0, W > 0) &= -N k_B T \ln 2 - \frac{1}{3} N k_B T e^{-W/k_B T} + \dots \\ S(T \rightarrow 0, W > 0) &= N \ln 2 \quad . \end{aligned}$$

Thus, the ground state entropies are the same, even though the allowed microstates are very different. For $W < 0$, there are no vacancies. For $W > 0$, every link must contain at least one vacancy.

(6.2) The Blume-Capel model is a spin-1 version of the Ising model, with Hamiltonian

$$H = -J \sum_{\langle ij \rangle} S_i S_j - \Delta \sum_i S_i^2 \quad ,$$

where $S_i \in \{-1, 0, +1\}$ and where the first sum is over all links of a lattice and the second sum is over all sites. It has been used to describe magnetic solids containing vacancies ($S = 0$ for a vacancy) as well as phase separation in $^4\text{He} - ^3\text{He}$ mixtures ($S = 0$ for a ^4He atom). For parts (b), (c), and (d) you should work in the thermodynamic limit. The eigenvalues and eigenvectors are such that it would shorten your effort considerably to use a program like Mathematica to obtain them.

- (a) Find the transfer matrix for the $d = 1$ Blume-Capel model.
- (b) Find the free energy $F(T, \Delta, N)$.
- (c) Find the density of $S = 0$ sites as a function of T and Δ .
- (d) *Exciting!* Find the correlation function $\langle S_j S_{j+n} \rangle$.

Solution :

(a) The transfer matrix R can be written in a number of ways, but it is aesthetically pleasing to choose it to be symmetric. In this case we have

$$R_{SS'} = e^{\beta J S S'} e^{\beta \Delta (S^2 + S'^2)/2} = \begin{pmatrix} e^{\beta(\Delta+J)} & e^{\beta\Delta/2} & e^{\beta(\Delta-J)} \\ e^{\beta\Delta/2} & 1 & e^{\beta\Delta/2} \\ e^{\beta(\Delta-J)} & e^{\beta\Delta/2} & e^{\beta(\Delta+J)} \end{pmatrix} \quad .$$

(b) For an N -site ring, we have

$$Z = \text{Tr} e^{-\beta H} = \text{Tr} (R^N) = \lambda_+^N + \lambda_0^N + \lambda_-^N \quad ,$$

where λ_+ , λ_0 , and λ_- are the eigenvalues of the transfer matrix R . To find the eigenvalues, note that

$$\vec{\psi}_0 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$$

is an eigenvector with eigenvalue $\lambda_0 = 2 e^{\beta\Delta} \sinh(\beta J)$. The remaining eigenvectors must be orthogonal to ψ_0 , and hence are of the form

$$\vec{\psi}_{\pm} = \frac{1}{\sqrt{2 + x_{\pm}^2}} \begin{pmatrix} 1 \\ x_{\pm} \\ 1 \end{pmatrix} \quad .$$

We now demand

$$R \begin{pmatrix} 1 \\ x \\ 1 \end{pmatrix} = \begin{pmatrix} 2 e^{\beta\Delta} \cosh(\beta J) + x e^{\beta\Delta/2} \\ 2 e^{\beta\Delta/2} + x \\ 2 e^{\beta\Delta} \cosh(\beta J) + x e^{\beta\Delta/2} \end{pmatrix} = \begin{pmatrix} \lambda \\ \lambda x \\ \lambda \end{pmatrix} \quad ,$$

resulting in the coupled equations

$$\begin{aligned} \lambda &= 2 e^{\beta\Delta} \cosh(\beta J) + x e^{\beta\Delta/2} \\ \lambda x &= 2 e^{\beta\Delta/2} + x \quad . \end{aligned}$$

Eliminating x , one obtains a quadratic equation for λ . The solutions are

$$\lambda_{\pm} = \left(e^{\beta\Delta} \cosh(\beta J) + \frac{1}{2} \right) \pm \sqrt{\left(e^{\beta\Delta} \cosh(\beta J) + \frac{1}{2} \right)^2 + 2 e^{\beta\Delta}}$$

$$x_{\pm} = e^{-\beta\Delta/2} \left\{ \left(\frac{1}{2} - e^{\beta\Delta} \cosh(\beta J) \right) \pm \sqrt{\left(\frac{1}{2} - e^{\beta\Delta} \cosh(\beta J) \right)^2 + 2 e^{\beta\Delta}} \right\} .$$

Note $\lambda_+ > \lambda_0 > 0 > \lambda_-$ and that λ_+ is the eigenvalue of the largest magnitude. This is in fact guaranteed by the *Perron-Frobenius theorem*, which states that for any positive matrix R (i.e. a matrix whose elements are all positive) there exists a positive real number p such that p is an eigenvalue of R and any other (possibly complex) eigenvalue of R is smaller than p in absolute value. Furthermore the associated eigenvector $\vec{\psi}$ is such that all its components are of the same sign. In the thermodynamic limit $N \rightarrow \infty$ we then have

$$F(T, \Delta, N) = -Nk_B T \ln \lambda_+ .$$

(c) Note that, at any site,

$$\langle S^2 \rangle = -\frac{1}{N} \frac{\partial F}{\partial \Delta} = \frac{1}{\beta} \frac{\partial \ln \lambda_+}{\partial \Delta} ,$$

and furthermore that

$$\delta_{S,0} = 1 - S^2 .$$

Thus,

$$\nu_0 \equiv \frac{N_0}{N} = 1 - \frac{1}{\beta} \frac{\partial \ln \lambda_+}{\partial \Delta} .$$

After some algebra, find

$$\nu_0 = 1 - \frac{r - \frac{1}{2}}{\sqrt{r^2 + 2 e^{\beta\Delta}}} ,$$

where

$$r = e^{\beta\Delta} \cosh(\beta J) + \frac{1}{2} .$$

It is now easy to explore the limiting cases $\Delta \rightarrow -\infty$, where we find $\nu_0 = 1$, and $\Delta \rightarrow +\infty$, where we find $\nu_0 = 0$. Both these limits make physical sense.

(d) We have

$$C(n) = \langle S_j S_{j+n} \rangle = \frac{\text{Tr}(\Sigma R^n \Sigma R^{N-n})}{\text{Tr}(R^N)} ,$$

where $\Sigma_{SS'} = S \delta_{SS'}$. We work in the thermodynamic limit. Note that $\langle + | \Sigma | + \rangle = 0$, therefore we must write

$$R = \lambda_+ |+\rangle\langle +| + \lambda_0 |0\rangle\langle 0| + \lambda_- |-\rangle\langle -| ,$$

and we are forced to choose the middle term for the n instances of R between the two Σ matrices. Thus,

$$C(n) = \left(\frac{\lambda_0}{\lambda_+} \right)^n |\langle + | \Sigma | 0 \rangle|^2 .$$

We define the correlation length ξ by

$$\xi = \frac{1}{\ln(\lambda_+/\lambda_0)} ,$$

in which case

$$C(n) = A e^{-|n|/\xi} ,$$

where now we generalize to positive and negative values of n , and where

$$A = |\langle + | \Sigma | 0 \rangle|^2 = \frac{1}{1 + \frac{1}{2} x_+^2} .$$

(6.3) DC Comics superhero Clusterman and his naughty dog Henry are shown in Fig. 1. Clusterman, as his name connotes, is a connected diagram, but the diagram for Henry contains some disconnected pieces.

- (a) Interpreting the diagrams as arising from the Mayer cluster expansion, compute the symmetry factor s_γ for Clusterman.
- (b) What is the *total* symmetry factor for Henry and his disconnected pieces? What would the answer be if, unfortunately, another disconnected piece of the same composition were to be found?
- (c) What is the lowest order virial coefficient to which Clusterman contributes?




Figure 1: Mayer expansion diagrams for Clusterman and his dog.

Solution :

First of all, this is really disgusting and you should all be ashamed that you had anything to do with this problem.

(a) Clusterman's head gives a factor of 6 because the upper three vertices can be permuted among themselves in any of $3! = 6$ ways. Each of his hands gives a factor of 2 because each hand can be rotated by π about its corresponding arm. The arms themselves can be interchanged, by rotating his shoulders by π about his body axis (Clusterman finds this invigorating). Finally, the analysis for the hands and arms applies just as well to the feet and legs, so we conclude

$$s_\gamma = 6 \cdot (2^2 \cdot 2)^2 = 3 \cdot 2^7 = 384 \quad .$$

Note that an arm cannot be exchanged with a leg, because the two lower vertices on Clusterman's torso are not equivalent. Plus, that would be a really mean thing to do to Clusterman.

(b) Henry himself has no symmetries. The little pieces each have $s_\Delta = 3!$, and moreover they can be exchanged, yielding another factor of 2. So the total symmetry factor for Henry plus disconnected pieces is $s_{\Delta\Delta} = 2! \cdot (3!)^2 = 72$. Were another little piece of the same...er...consistency to be found, the symmetry factor would be $s_{\Delta\Delta\Delta} = 3! \cdot (3!)^3 = 2^4 \cdot 3^4 = 1296$, since we get a factor of $3!$ from each of the Δ pieces, and a fourth factor of $3!$ from the permutations among the Δ s.

(c) There are 18 vertices in Clusterman, hence he will first appear in B_{18} .

(6.4) Use the high temperature expansion to derive the spin-spin correlation functions for a spin- $\frac{1}{2}$ ($\sigma_n = \pm 1$) Ising chain and Ising ring. Compare with the results in chapter 6 of the lecture notes.

Solution :

The spin-spin correlation function $C_{kl} = \langle \sigma_k \sigma_l \rangle$ is expressed as a ratio Y_{kl}/Z as in eqn. 6.51 of the Lecture Notes (LN). For the chain, the only diagram which contributes to Z is $\Gamma = \{\emptyset\}$, i.e. the trivial empty lattice. This is because there is no way to form closed loops on a chain. Thus $Z_{\text{ring}} = 2^N (\cosh \beta J)^{N-1}$ since the number of links is $N_L = N - 1$ (see LN eqn. 6.45). For the chain, in addition to the empty lattice, there is one closed loop that can be formed which includes every link of the chain. Thus $Z_{\text{chain}} = 2^N (\cosh \beta J)^N (1 + x^N)$, where $x = \tanh \beta J$. As for the numerator Y_{kl} , on the chain there is only one possible string, shown in Fig. 2, which extends between sites k and l . Thus $Y_{kl}^{\text{chain}} = 2^N (\cosh \beta J)^{N-1} x^{|k-l|}$. On the ring there are two possible strings, since the ring is multiply connected. Thus $Y_{kl}^{\text{ring}} = 2^N (\cosh \beta J)^N (x^{|k-l|} + x^{N-|k-l|})$. Therefore,

$$C_{kl}^{\text{chain}} = x^{|k-l|} \quad , \quad C_{kl}^{\text{ring}} = \frac{x^{|k-l|} + x^{N-|k-l|}}{1 + x^N} \quad .$$




Figure 2: Diagrams for the numerator of the high temperature expansion of the spin-spin correlation function on an Ising ring and chain.

(6.5) An ionic solution of dielectric constant ϵ and mean ionic density n fills a grounded conducting sphere of radius R . A charge Q lies at the center of the sphere. Calculate the ionic charge density as a function of the radial coordinate r , assuming $Q/r \ll k_B T$.

Solution :

Debye-Hückel theory tells us that

$$n_{\pm}(\mathbf{r}) = \frac{1}{2}n_{\infty} e^{\mp e\phi(\mathbf{r})/k_B T}$$

and

$$\nabla^2 \phi = -\frac{4\pi e}{\epsilon} (n_+ - n_-) - \frac{4\pi}{\epsilon} \rho_{\text{ext}} \quad ,$$

where ϵ is the dielectric constant. Assuming $\phi \ll k_B T$, we have $\nabla^2 \phi = \kappa_D^2 \phi - 4\pi\epsilon^{-1}\rho_{\text{ext}}$, with

$$\kappa_D = \sqrt{\frac{4\pi n_{\infty} e^2}{\epsilon k_B T}} \quad .$$

Assuming a spherically symmetric solution, with a point charge Q at the origin, we solve

$$\left(-\frac{1}{r} \frac{\partial^2}{\partial r^2} r + \kappa_D^2 \right) \phi = \frac{4\pi Q}{\epsilon} \delta(\mathbf{r}) \quad .$$

The solution is then of the form $\phi(r) = \frac{1}{r} u(r)$, with $u'' = \kappa_D^2 u$ for $r > 0$. Thus,

$$\phi(r) = A \frac{\cosh(\kappa_D r)}{r} + B \frac{\sinh(\kappa_D r)}{r} \quad .$$

As $r \rightarrow 0$ we must have an unscreened charge Q , hence $A = Q/\epsilon$. The boundary condition on the conducting sphere is $\phi(R) = 0$, hence $B = -A \coth(\kappa_D R)$. Thus,

$$\phi(\mathbf{r}) = \frac{Q \cosh(\kappa_D r)}{\epsilon r} \cdot \left(1 - \frac{\tanh(\kappa_D r)}{\tanh(\kappa_D R)} \right) \quad .$$

We stress that this solution is valid only where $e\phi(\mathbf{r}) \ll k_B T$.

(6.6) Consider a three-dimensional gas of point particles interacting according to the potential

$$u(r) = \begin{cases} +\Delta_0 & \text{if } r \leq a \\ -\Delta_1 & \text{if } a < r \leq b \\ 0 & \text{if } b < r \end{cases},$$

where $\Delta_{0,1}$ are both positive. Compute the second virial coefficient $B_2(T)$ and find a relation which determines the inversion temperature in a throttling process.

Solution :

The Mayer function is

$$f(r) = \begin{cases} e^{-\Delta_0/k_B T} - 1 & \text{if } r \leq a \\ e^{\Delta_1/k_B T} - 1 & \text{if } a < r \leq b \\ 0 & \text{if } b < r \end{cases}.$$

The second virial coefficient is

$$\begin{aligned} B_2(T) &= -\frac{1}{2} \int d^3r f(r) \\ &= \frac{2\pi a^3}{3} \cdot \left[(1 - e^{-\Delta_0/k_B T}) + (s^3 - 1)(1 - e^{\Delta_1/k_B T}) \right], \end{aligned}$$

where $s = b/a$. The inversion temperature is a solution of the equation $B_2(T) = TB_2'(T)$, which gives

$$s^3 - 1 = \frac{1 + \left(\frac{\Delta_0}{k_B T} - 1\right) e^{-\Delta_0/k_B T}}{1 + \left(\frac{\Delta_1}{k_B T} + 1\right) e^{\Delta_1/k_B T}}.$$

(6.7) At the surface of every metal a dipolar layer develops which lowers the potential energy for electrons inside the metal. Some electrons near the surface escape to the outside, leaving a positively charged layer behind, while overall there is charge neutrality. The situation is depicted in Fig. 3. The electron density outside the metal is very low and Maxwell-Boltzmann statistics are appropriate.

(a) Consider a flat metallic surface, normal to \hat{x} , located at $x = 0$. Assume for $x > 0$ an electronic distribution $n(x) = n_0 \exp(e\phi/k_B T)$, where ϕ is the electric potential. For $x > 0$ there are only electrons; all the positive charges are located within the metal. Write down the self-consistent equation for the potential $\phi(x)$.

(b) Having found the self-consistent equation for $\phi(x)$, show that, multiplying by $\phi'(x)$, the equation can be integrated once, analogous to the conservation of energy for mechanical systems (with ϕ playing the role of the coordinate and x playing the role of time). Show that the equation can be integrated once again to yield $\phi(x)$, with the constant determined by the requirement that $n(x=0) = n_0$.

(c) Find $n(x)$.

Solution :

(a) The self-consistent equation is Poisson's equation,

$$\nabla^2 \phi = -4\pi\rho = 4\pi en_0 e^{e\phi/k_B T} .$$

Since the only variation is along x , we have $\phi'' = 4\pi en_0 e^{e\phi/k_B T}$. Multiplying each side by $\frac{d\phi}{dx}$, we have

$$\frac{d}{dx} \left(\frac{1}{2} \phi'^2 \right) = \frac{d}{dx} \left(4\pi n_0 k_B T e^{e\phi/k_B T} \right) ,$$

and integrating this equation from x to ∞ we obtain

$$\frac{d\phi}{dx} = -(8\pi n_0 k_B T)^{1/2} e^{e\phi/2k_B T} .$$

Note also the choice of sign here, due to the fact that the potential $-e\phi$ for electrons must increase with x . The boundary term at $x = \infty$ must vanish since $n(\infty) = 0$, which requires $e^{e\phi(\infty)/k_B T} = 0$.

(b) Integrating once more, we have

$$e^{-e\phi(x)/2k_B T} = \left(\frac{2\pi n_0 e^2}{k_B T} \right)^{1/2} (x + a) ,$$

where a is a constant of integration. Since $n(x=0) \equiv n_0$, we must have $\phi(0) = 0$, and hence

$$a = \left(\frac{k_B T}{2\pi n_0 e^2} \right)^{1/2} .$$

Thus,

$$\phi(x) = -\frac{2k_B T}{e} \ln \left(\frac{x+a}{a} \right) .$$

(c) The electron number distribution is then

$$n(x) = n_0 \left(\frac{a}{x+a} \right)^2 .$$




Figure 3: Electron distribution in the vicinity of the surface of a metal.

(6.8) In §6.4.3 of the notes, the virial equation of state is derived for a single species of particle.

- (a) Generalize eqn. 5.160 to the case of two species interacting by $u_{\sigma\sigma'}(r)$, where σ and σ' are the species labels.
- (b) For a plasma, show from Debye-Hückel theory that the pair correlation function is $g_{\sigma\sigma'} \propto \exp(-\sigma\sigma'q^2\phi(r)/k_B T)$, where σ and σ' are the signs of the charges (magnitude q), and $\phi(r)$ is the screened potential due to a unit positive test charge.
- (c) Find the equation of state for a three-dimensional two-component plasma, in the limit where T is large.

Solution :

(a) Let $i = 1, \dots, N_+ + N_-$ index all the particles, and let $\sigma_i = \pm 1$ denote the sign of the charge of particle i , with $\sigma_i = +1$ for $1 \leq i \leq N_+$ and $\sigma_i = -1$ for $(N_+ + 1) \leq i \leq (N_+ + N_-)$. In a globally neutral system, $N_+ = N_- \equiv \frac{1}{2}N$. We define

$$g_{\mu\nu}(\mathbf{r}) \equiv \frac{1}{n_\mu n_\nu} \left\langle \sum_{i \neq j} \delta(\mathbf{r} - \mathbf{x}_i) \delta(\mathbf{x}_j) \delta_{\sigma_i, \mu} \delta_{\sigma_j, \nu} \right\rangle ,$$

where n_μ is the density of particles of species μ , with $\mu = \pm 1$. As defined, $g_{\mu\nu}(\mathbf{r}) \rightarrow 1$ as $r \rightarrow \infty$. If instead we normalize $g_{\mu\nu}$ by dividing by $n_{\text{tot}}^2 = (n_+ + n_-)^2$, then we would have $g_{\mu\nu}(r \rightarrow \infty) = \frac{1}{4}$. We next work on the virial equation of state,

$$\frac{p}{k_B T} = \frac{N_+ + N_-}{V} - \frac{1}{3V k_B T} \sum_{i=1}^{N_+ + N_-} \langle \mathbf{x}_i \cdot \nabla_i W \rangle .$$

The potential is

$$W = \sum_{i < j} \frac{\sigma_i \sigma_j q^2}{|\mathbf{x}_i - \mathbf{x}_j|} \equiv \sum_{i < j} u_{\sigma_i \sigma_j}(|\mathbf{x}_i - \mathbf{x}_j|) ,$$

with $u_{\sigma\sigma'}(r) = \sigma\sigma'q^2/r$. Then using translational invariance one has

$$\frac{p}{k_B T} = n_+ + n_- - \frac{2\pi}{3k_B T} \sum_{\sigma, \sigma'} n_\sigma n_{\sigma'} \int_0^\infty dr r^3 u'_{\sigma\sigma'}(r) g_{\sigma\sigma'}(r)$$

(b) According to Debye-Hückel theory,

$$g_{\sigma\sigma'}(r) = \exp\left(-\frac{\sigma\sigma'q\phi(r)}{k_B T}\right) ,$$

where $\phi(r)$ is the screened potential at r due to a point charge q at the origin, which satisfies

$$\nabla^2 \phi = 4\pi n q \sinh(q\phi/k_B T) - 4\pi q \delta(\mathbf{r}) ,$$

where $n_+ = n_- \equiv \frac{1}{2}n$. In the high temperature limit, we can expand the sinh function and we obtain the Yukawa potential

$$\phi(r) = \frac{q}{r} e^{-\kappa_D r} ,$$

where

$$\kappa_D = \left(\frac{4\pi n q^2}{k_B T}\right)^{1/2}$$

is the Debye screening wavevector. Thus, we have

$$\begin{aligned}
\frac{p}{k_{\text{B}}T} &= n - \frac{\pi n^2}{6k_{\text{B}}T} \int_0^\infty dr r^3 \left(-\frac{q^2}{r^2} \right) \sum_{\sigma, \sigma'} \sigma \sigma' g_{\sigma \sigma'}(r) \\
&= n - \frac{2\pi n^2 q^3}{3(k_{\text{B}}T)^2} \int_0^\infty dr r \phi(r) = n - \frac{2\pi n^2 q^4}{3(k_{\text{B}}T)^2 \kappa_{\text{D}}} \\
&= n \left(1 - \frac{\sqrt{\pi} n^{1/2} q^3}{3(k_{\text{B}}T)^{3/2}} \right) .
\end{aligned}$$

(6.9) Consider a liquid where the interaction potential is $u(r) = \Delta_0 (a/r)^k$, where Δ_0 and a are energy and length scales, respectively. Assume that the pair distribution function is given by $g(r) \approx e^{-u(r)/k_B T}$. Compute the equation of state. For what values of k do your expressions converge?

Solution:

According to the virial equation of state in eqn. 6.157 of the Lecture Notes,

$$p = nk_B T - \frac{2}{3} \pi n^2 \int_0^{\infty} dr r^3 g(r) u'(r) \quad .$$

Substituting for $u(r)$ and $g(r)$ as in the statement of the problem, we change variables to

$$s \equiv \frac{u(r)}{k_B T} \quad \Rightarrow \quad ds = \frac{u'(r)}{k_B T} dr \quad ,$$

so

$$r = a \left(\frac{\Delta_0}{k_B T} \right)^{1/k} s^{-1/k}$$

and

$$r^3 g(r) u'(r) dr = k_B T a^3 \left(\frac{\Delta_0}{k_B T} \right)^{3/k} s^{-3/k} e^{-s} ds \quad .$$

We then have

$$\begin{aligned} p &= nk_B T + \frac{2}{3} \pi n^3 a^3 k_B T \left(\frac{\Delta_0}{k_B T} \right)^{3/k} \int_0^{\infty} ds s^{-3/k} e^{-s} \\ &= nk_B T \left\{ 1 + \frac{2}{3} \pi \Gamma \left(1 - \frac{3}{k} \right) n a^3 \left(\frac{\Delta_0}{k_B T} \right)^{3/k} \right\} \quad . \end{aligned}$$

Note that a minus sign appears because we must switch the upper and lower limits on the s integral. This expression converges provided $k < 0$ or $k > 3$.

(6.10) Consider a charge Q impurity located at the origin of a two-dimensional metallic plane. You may model the plane initially as a noninteracting Fermi gas in the presence of a neutralizing background. Poisson's equation is

$$\nabla^2 \phi = 4\pi e [n(\boldsymbol{\rho}) - n_0] \delta(z) - 4\pi Q \delta(\boldsymbol{\rho}) \delta(z) \quad ,$$

where $\mathbf{r} = (\boldsymbol{\rho}, z)$ is decomposed into a two-dimensional vector $\boldsymbol{\rho}$ and the scalar z , and where n_0 is the number density of electrons at $|\boldsymbol{\rho}| = \infty$.

- (a) Using the Thomas-Fermi approach, find the two-dimensional electron number density $n(\boldsymbol{\rho})$ in terms of the local potential $\phi(\boldsymbol{\rho}, 0)$.
- (b) By Fourier transformation, show that

$$\hat{\phi}(\mathbf{k}, q) = \frac{4\pi Q}{\mathbf{k}^2 + q^2} - \frac{4\pi n_0 e^2}{\varepsilon_F} \frac{\hat{\chi}(\mathbf{k})}{\mathbf{k}^2 + q^2} \quad ,$$

where \mathbf{k} is a two-dimensional wavevector, and

$$\hat{\chi}(\mathbf{k}) = \int_{-\infty}^{\infty} \frac{dq}{2\pi} \hat{\phi}(\mathbf{k}, q) \quad .$$

- (c) Solve for $\hat{\chi}(\mathbf{k})$ and then for $\hat{\phi}(\mathbf{k}, q)$.
- (d) Derive an expression for the potential $\phi(\boldsymbol{\rho}, z)$.
- (e) Derive an expression for the local charge density $\varrho(\boldsymbol{\rho}) = en_0 - en(\boldsymbol{\rho})$. Show that $\varrho(\boldsymbol{\rho}) = \frac{Q}{2\pi\lambda^2} f(\rho/\lambda)$, where λ is a screening length and $f(s)$ is some function, and expression for which you should derive. Sketch $f(s)$.

Solution:

(a) In two dimensions we have

$$n = 2 \int \frac{d^2k}{(2\pi)^2} \Theta(k_F - k) = \frac{k_F^2}{2\pi} = \frac{m\varepsilon_F}{\pi\hbar^2} \quad ,$$

where we have used $\varepsilon_F = \hbar^2 k_F^2 / 2m$. In the presence of a potential, the energy levels are shifted and it is the electrochemical potential $\varepsilon_F^\infty = \varepsilon_F - e\phi$ which is constant throughout the system. Thus, the local electron density is

$$n(\boldsymbol{\rho}) = \frac{m}{\pi\hbar^2} \left[\varepsilon_F^\infty + e\phi(\boldsymbol{\rho}, 0) \right] = n_0 + \frac{me}{\pi\hbar^2} \phi(\boldsymbol{\rho}, 0) \quad .$$

Here, $\phi(\mathbf{r}) = \phi(\boldsymbol{\rho}, z)$ is the electrostatic potential in three-dimensional space. When we restrict to the $z = 0$ plane we write $\phi(\boldsymbol{\rho}, 0)$.

(b) We now have

$$\nabla^2 \phi = \frac{4}{a_B} \phi(\boldsymbol{\rho}, 0) \delta(z) - 4\pi Q \delta(\boldsymbol{\rho}) \delta(z) \quad ,$$

where $a_B = \hbar^2 / me^2$ is the Bohr radius. Now we take the Fourier transform by multiplying the above equation by $e^{i\mathbf{k}\cdot\boldsymbol{\rho}} e^{iqz}$ and then integrating over all $\boldsymbol{\rho}$ and z . This gives

$$-(\mathbf{k}^2 + q^2) \hat{\phi}(\mathbf{k}, q) = \frac{4}{a_B} \overbrace{\int_{-\infty}^{\infty} \frac{dq}{2\pi} \hat{\phi}(\mathbf{k}, q)}^{\hat{\chi}(\mathbf{k})} - 4\pi Q \quad ,$$

```
Integrate[BesselJ[0, u a] / (1 + u), {u, 0, Infinity}, Assumptions -> Re[a] > 0 && Abs[Im[a]] == 0]
```

$$\frac{1}{2} \pi (-\text{BesselY}[0, a] + \text{StruveH}[0, a])$$

```
F[x_] := 1 / x + \frac{1}{2} \pi (\text{BesselY}[0, x] - \text{StruveH}[0, x])
```

```
Plot[F[x], {x, 0, 10}, AxesLabel -> {\sigma, F[\sigma, 0]}, PlotStyle -> Thick]
```




Figure 4: Plot of the screening charge density in units of $-Q/2\pi\lambda^2$ for problem (10).

hence

$$\hat{\phi}(\mathbf{k}, q) = \frac{4\pi Q}{\mathbf{k}^2 + q^2} - \frac{4}{a_B} \frac{\hat{\chi}(\mathbf{k})}{\mathbf{k}^2 + q^2} \quad .$$

(c) To solve for $\hat{\chi}(\mathbf{k})$ we integrate the above equation over q and use the fact that

$$\int_{-\infty}^{\infty} \frac{dq}{2\pi} \frac{e^{iqz}}{\mathbf{k}^2 + q^2} = \frac{e^{-|kz|}}{2|\mathbf{k}|} \quad .$$

Thus,

$$\hat{\chi}(\mathbf{k}) = \frac{2\pi Q}{|\mathbf{k}|} - \frac{2}{|ka_B|} \hat{\chi}(\mathbf{k})$$

Thus,

$$\hat{\chi}(\mathbf{k}) = \frac{2\pi Q}{|\mathbf{k}| + \lambda^{-1}} \quad ,$$

where $\lambda = \frac{1}{2}a_B$. Plugging this back into our equation for $\hat{\phi}(\mathbf{k}, q)$, we obtain

$$\hat{\phi}(\mathbf{k}, q) = \frac{4\pi Q \cdot |\mathbf{k}\lambda|}{(\mathbf{k}^2 + q^2)(1 + |\mathbf{k}\lambda|)} \quad .$$

(d) Now we Fourier transform back to real space:

$$\begin{aligned}
\phi(\boldsymbol{\rho}, z) &= \int \frac{d^2k}{(2\pi)^2} \int_{-\infty}^{\infty} \frac{dq}{2\pi} \hat{\phi}(\mathbf{k}, q) e^{i\mathbf{k}\cdot\boldsymbol{\rho}} e^{iqz} \\
&= \int \frac{d^2k}{(2\pi)^2} \frac{e^{-|kz|}}{2|\mathbf{k}|} \cdot \frac{4\pi Q |\mathbf{k}\lambda|}{1 + |\mathbf{k}\lambda|} \cdot e^{i\mathbf{k}\cdot\boldsymbol{\rho}} \\
&= \frac{Q}{\lambda} F(\rho/\lambda, |z|/\lambda) \quad ,
\end{aligned}$$

where

$$F(\sigma, \zeta) = \int_0^{\infty} du \frac{u}{1+u} J_0(\sigma u) e^{-\zeta u} \quad ,$$

where $J_0(s)$ is the Bessel function of order zero.

(e) We have

$$\varrho(\boldsymbol{\rho}) = e[n_0 - n(\boldsymbol{\rho})] = -\frac{Q}{2\pi\lambda^2} F(\rho/\lambda, 0) \quad .$$

Note

$$\begin{aligned}
F(\rho/\lambda, 0) &= \int_0^{\infty} du \frac{u J_0(u\rho/\lambda)}{1+u} = \frac{\lambda}{\rho} - \int_0^{\infty} du \frac{J_0(u\rho/\lambda)}{1+u} \\
&= \frac{\lambda}{\rho} + \frac{1}{2}\pi Y_0(\rho/\lambda) - \frac{1}{2}\pi \mathbf{H}_0(\rho/\lambda) \quad ,
\end{aligned}$$

where $Y_0(s)$ is a Bessel function of the second kind and $\mathbf{H}_0(s)$ is the Struve function. Asymptotically¹ we obtain

$$\varrho(\boldsymbol{\rho}) = \frac{Q}{2\pi\lambda^2} \left\{ \sum_{n=1}^{p-1} (-1)^n \Gamma^2\left(\frac{1}{2} + n\right) \left(\frac{2\lambda}{\rho}\right)^{(2n+1)} + \mathcal{O}(2\lambda/\rho)^{2p+1} \right\} \quad .$$

Note that $\varrho(\rho) \propto \rho^{-3}$ at large distances. In the above formula, p is arbitrary. Since $\Gamma(z + \frac{1}{2}) \sim z \ln z - z$, the optimal value of p to minimize the remainder in the sum is $p \approx \rho/2\lambda$. See Fig. 4 for a sketch.

¹See Gradshteyn and Ryzhik §8.554, then use $\Gamma(z)\Gamma(1-z) = \pi \csc(\pi z)$.

(11) The grand partition function for a system is given by the expression

$$\Xi = (1 + z)^{V/v_0} (1 + z^{\alpha V/v_0}) \quad ,$$

where $\alpha > 0$. In this problem, you are to work in the thermodynamic limit. You will also need to be careful to distinguish the cases $|z| < 1$ and $|z| > 1$.

- (a) Find an expression for the pressure $p(T, z)$.
- (b) Find an expression for the number density $n(T, z)$.
- (c) Plot $v(p, T)$ as a function of p for different temperatures and show there is a first order phase transition, *i.e.* a discontinuity in $v(p)$, which occurs for $|z| = 1$. What is the change in volume at the transition? .

Solution :

(a) The grand potential is

$$\Omega(T, z) = -k_B T \ln \Xi = -\frac{k_B T V}{v_0} \ln(1 + z) - k_B T \ln(1 + z^{\alpha V/v_0}) \quad .$$

Now take the thermodynamic limit $V/v_0 \rightarrow \infty$. One then has

$$\Omega(T, z) = -\frac{k_B T V}{v_0} \ln(1 + z) - \begin{cases} 0 & \text{if } |z| < 1 \\ \frac{\alpha k_B T V}{v_0} \ln z & \text{if } |z| > 1 \end{cases} \quad .$$

From this we compute the pressure,

$$\begin{aligned} p &= -\left(\frac{\partial \Omega}{\partial V}\right)_{T, \mu} = \frac{k_B T}{v_0} \ln(1 + z) + \frac{\alpha k_B T}{v_0} \cdot \frac{z^{\alpha V/v_0} \ln z}{1 + z^{\alpha V/v_0}} \\ &= \frac{k_B T}{v_0} \ln(1 + z) + \begin{cases} 0 & \text{if } |z| < 1 \\ \frac{\alpha k_B T}{v_0} \ln z & \text{if } |z| > 1 \end{cases} \quad . \end{aligned}$$

(b) For the density, we have

$$\begin{aligned} n &= -\frac{z}{V k_B T} \left(\frac{\partial \Omega}{\partial z}\right)_{T, V} = \frac{1}{v_0} \cdot \frac{z}{1 + z} + \frac{\alpha}{v_0} \cdot \frac{z^{\alpha V/v_0}}{1 + z^{\alpha V/v_0}} \\ &= \frac{1}{v_0} \cdot \frac{z}{1 + z} + \begin{cases} 0 & \text{if } |z| < 1 \\ \alpha/v_0 & \text{if } |z| > 1 \end{cases} \quad . \end{aligned}$$

(c) We eliminate z from the above equations, and we write $v = 1/n$ as the volume per particle. The fugacity $z(v)$ satisfies

$$z(v) = \begin{cases} \frac{v_0}{v - v_0} & \text{if } v > 2v_0 \\ 1 & \text{if } \frac{2v_0}{1 + 2\alpha} < v < 2v_0 \\ \frac{v_0 - \alpha v}{(1 + \alpha)v - v_0} & \text{if } \frac{v_0}{1 + \alpha} < v < \frac{2v_0}{1 + 2\alpha} \\ \infty & \text{if } v < \frac{v_0}{1 + \alpha} \end{cases}$$

We then have

$$\frac{pv_0}{k_B T} = \begin{cases} \ln\left(\frac{v}{v-v_0}\right) & v > 2v_0 \\ \ln 2 & \frac{2v_0}{1+2\alpha} < v < 2v_0 \\ \ln\left[\left(\frac{v}{(1+\alpha)v-v_0}\right)\left(\frac{v_0-\alpha v}{(1+\alpha)v-v_0}\right)^\alpha\right] & \frac{v_0}{1+\alpha} < v < \frac{2v_0}{1+2\alpha} \\ \infty & v < \frac{v_0}{1+\alpha} \end{cases}$$

Sample plots of $z(v)$ and $p(v)$ are shown in Fig. 5.




Figure 5: $z(v)$ and $p(v)$ for $\alpha = 0.2, 1.0$, and 3.0 .

(6.12) In problem 11, you considered the thermodynamic properties associated with the grand partition function $\Xi(V, z) = (1+z)^{V/v_0} (1+z^{\alpha V/v_0})$. Consider now the following partition function:

$$\Xi(V, z) = (1+z)^{V/v_0} \prod_{j=1}^K \left\{ 1 + \left(\frac{z}{\sigma_j} \right)^{\alpha V/Kv_0} \right\} .$$

Consider the thermodynamic limit where α is a number on the order of unity, $V/v_0 \rightarrow \infty$, and $K \rightarrow \infty$ but with $Kv_0/V \rightarrow 0$. For example, we might have $K \propto (V/v_0)^{1/2}$.

(a) Show that the number density is

$$n(T, z) = \frac{1}{v_0} \frac{z}{1+z} + \frac{\alpha}{v_0} \int_0^{|z|} d\sigma g(\sigma) ,$$

where

$$g(\sigma) = \frac{1}{K} \sum_{j=1}^K \delta(\sigma - \sigma_j) .$$

(b) Derive the corresponding expression for $p(T, z)$.

(c) In the thermodynamic limit, the spacing between consecutive σ_j values becomes infinitesimal. In this case, $g(\sigma)$ approaches a continuous distribution. Consider the flat distribution,

$$g(\sigma) = \frac{1}{w} \Theta(\sigma - r) \Theta(r + w - \sigma) = \begin{cases} w^{-1} & \text{if } r < \sigma < r + w \\ 0 & \text{otherwise.} \end{cases}$$

The model now involves three dimensionless parameters²: α , r , and w . Solve for $z(v)$. You will have to take cases, and you should find there are three regimes to consider³.

(d) Plot $pv_0/k_B T$ versus v/v_0 for the case $\alpha = \frac{1}{4}$ and $r = w = 1$.

(e) Comment on the critical properties (*i.e.* the singularities) of the equation of state.

Solution :

(a) We have

$$\frac{1}{V} \ln \Xi = \frac{1}{v_0} \ln(1+z) + \frac{\alpha}{Kv_0} \sum_{i=1}^K \ln(z/\sigma_i) \Theta(|z| - \sigma_i) ,$$

so from $n = V^{-1} z \partial \ln \Xi / \partial z$,

$$\begin{aligned} n &= \frac{1}{v_0} \frac{z}{1+z} + \frac{\alpha}{Kv_0} \sum_{i=1}^K \Theta(|z| - \sigma_i) \\ &= \frac{1}{v_0} \frac{z}{1+z} + \frac{\alpha}{v_0} \int_0^{|z|} d\sigma g(\sigma) . \end{aligned}$$

²The quantity v_0 has dimensions of volume and disappears from the problem if one defines $\tilde{v} = v/v_0$.

³You should find that a fourth regime, $v < (1+r^{-1})v_0$, is not permitted.

(b) The pressure is $p = V^{-1}k_B T \ln \Xi$:

$$\begin{aligned} p &= \frac{k_B T}{v_0} \ln(1+z) + \frac{\alpha k_B T}{K v_0} \sum_{i=1}^K \ln(z/\sigma_i) \Theta(|z| - \sigma_i) \\ &= \frac{k_B T}{v_0} \ln(1+z) + \frac{\alpha k_B T}{v_0} \int_0^{|z|} d\sigma g(\sigma) \ln(z/\sigma) \quad . \end{aligned}$$

(c) We now consider the given form for $g(\sigma)$. From our equation for $n(z)$, we have

$$n v_0 = \frac{v_0}{v} = \begin{cases} \frac{z}{1+z} & \text{if } |z| \leq r \\ \frac{z}{1+z} + \frac{\alpha}{w} (z-r) & \text{if } r \leq |z| \leq r+w \\ \frac{z}{1+z} + \alpha & \text{if } r+w \leq |z| \quad . \end{cases}$$

We need to invert this result. We assume $z \in \mathbb{R}^+$. In the first regime, we have

$$z \in [0, r] \quad \Rightarrow \quad z = \frac{v_0}{v - v_0} \quad \text{with} \quad \frac{v}{v_0} \in [1 + r^{-1}, \infty] \quad .$$

In the third regime,

$$z \in [r+w, \infty] \quad \Rightarrow \quad z = \frac{v_0 - \alpha v}{(1+\alpha)v - v_0} \quad \text{with} \quad \frac{v}{v_0} \in \left[\frac{1}{1+\alpha}, \frac{1+r+w}{(1+\alpha)(r+w)+\alpha} \right] \quad .$$

Note that there is a minimum possible volume per particle, $v_{\min} = v_0/(1+\alpha)$, hence a maximum possible density $n_{\max} = 1/v_{\min}$. This leaves us with the second regime, where $z \in [r, r+w]$. We must invert the relation

$$\frac{v_0}{v} = \frac{z}{1+z} + \frac{\alpha}{w} (z-r) \quad \Rightarrow \quad \frac{\alpha}{w} z^2 + \left(\frac{\alpha}{w} (1-r) + 1 - \frac{v_0}{v} \right) z - \left(\frac{\alpha r}{w} + \frac{v_0}{v} \right) = 0 \quad .$$

obtaining

$$z = \frac{-\left[\frac{\alpha}{w} (1-r) + 1 - \frac{v_0}{v} \right] + \sqrt{\left[\frac{\alpha}{w} (1-r) + 1 - \frac{v_0}{v} \right]^2 + \frac{4\alpha}{w} \left(\frac{\alpha r}{w} + \frac{v_0}{v} \right)}}{2\alpha/w} \quad ,$$

which holds for

$$a \in [r, r+w] \quad \Rightarrow \quad \frac{v}{v_0} \in \left[\frac{1+r+w}{(1+\alpha)(r+w)+\alpha}, 1+r^{-1} \right] \quad .$$

The dimensionless pressure $\pi = p v_0 / k_B T$ is given by

$$z \in [0, r] \quad \Rightarrow \quad \pi = \ln(1+z) \quad \text{with} \quad \frac{v}{v_0} \in [1 + r^{-1}, \infty] \quad .$$

and

$$z \in [r+w, \infty] \quad \Rightarrow \quad \pi = \ln(1+z) + \alpha \ln z - \frac{\alpha}{w} \left[(r+w) \ln(r+w) - r \ln r - w \right]$$

in the large volume region and

$$\frac{v}{v_0} \in \left[\frac{1}{1+\alpha}, \frac{1+r+w}{(1+\alpha)(r+w)+\alpha} \right]$$

in the small volume region. In the intermediate volume region, we have

$$\pi = \ln(1+z) + \frac{\alpha}{w} (z-r) \ln z - \frac{\alpha}{w} (z \ln z - r \ln r - z + r) \quad ,$$

which holds for

$$z \in [r, r+w] \Rightarrow \frac{v}{v_0} \in \left[\frac{1+r+w}{(1+\alpha)(r+w)+\alpha}, 1+r^{-1} \right] .$$

(d) The results are plotted in Fig. 4. Note that v is a continuous function of π , indicating a second order transition.

(e) Consider the thermodynamic behavior in the vicinity of $z = r$, *i.e.* near $v = (1+r^{-1})v_0$. Let's write $z = r + \epsilon$ and work to lowest nontrivial order in ϵ . On the low density side of this transition, *i.e.* for $\epsilon < 0$, we have, with $\nu = nv_0 = v_0/v$,

$$\nu = \frac{z}{1+z} = \frac{r}{1+r} + \frac{\epsilon}{(1+r)^2} + \mathcal{O}(\epsilon^2)$$

$$\pi = \ln(1+z) = \ln(1+r) + \frac{\epsilon}{1+r} + \mathcal{O}(\epsilon^2) .$$

Eliminating ϵ , we have

$$\nu < \nu_c \Rightarrow \pi = \ln(1+r) + (1+r)(\nu - \nu_c) + \dots ,$$

where $\nu_c = r/(1+r)$ is the critical dimensionless density. Now investigate the high density side of the transition, where $\epsilon > 0$. Integrating over the region $[r, r+\epsilon]$, we find

$$\nu = \frac{z}{1+z} + \frac{\alpha}{w}(z-r) = \frac{r}{1+r} + \left[\frac{1}{(1+r)^2} + \frac{\alpha}{w} \right] \epsilon + \mathcal{O}(\epsilon^2)$$

$$\pi = \ln(1+z) + \frac{\alpha}{w} [z + r \ln(r/z) - r] = \ln(1+r) + \frac{\epsilon}{1+r} + \mathcal{O}(\epsilon^2) .$$




Figure 6: Fugacity z and dimensionless pressure $pv_0/k_B T$ versus dimensionless volume per particle v/v_0 for problem (2), with $\alpha = \frac{1}{4}$ and $r = w = 1$. Different portions of the curves are shown in different colors. The dashed line denotes the minimum possible volume $v_{\min} = v_0/(1+\alpha)$.

Note that $\partial\pi/\partial z$ is continuous through the transition. As we are about to discover, $\partial\pi/\partial\nu$ is discontinuous. Eliminating ϵ , we have

$$\nu > \nu_c \quad \Rightarrow \quad \pi = \ln(1+r) + \frac{1+r}{1+(1+r)^2(\alpha/w)}(\nu - \nu_c) + \dots \quad .$$

Thus, the isothermal compressibility $\kappa_T = -\frac{1}{\nu}(\frac{\partial\nu}{\partial p})_T$ is discontinuous at the transition. This can be seen clearly as a kink in Fig. 6.

Suppose the density of states $g(\sigma)$ behaves as a power law in the vicinity of $\sigma = r$, with $g(\sigma) \simeq A(\sigma - r)^t$. Normalization of the integral of $g(\sigma)$ then requires $t > -1$ for convergence at this lower limit. For $z = r + \epsilon$ with $\epsilon > 0$, one now has

$$\begin{aligned} \nu &= \frac{r}{1+r} + \frac{\epsilon}{(1+r)^2} + \frac{\alpha A \epsilon^{t+1}}{t+1} + \dots \\ \pi &= \ln(1+r) + \frac{\epsilon}{1+r} + \frac{\alpha A \epsilon^{t+2}}{(t+1)(t+2)r} + \dots \quad . \end{aligned}$$

If $t > 0$, then to order ϵ the expansion is the same for $\epsilon < 0$, and both π and its derivative $\frac{\partial\pi}{\partial\nu}$ are continuous across the transition. (Higher order derivatives, however, may be discontinuous or diverge.) If $-1 < t < 0$, then ϵ^{t+1} dominates over ϵ in the first of these equations, and we have

$$\epsilon = \left(\frac{(t+1)(\nu - \nu_c)}{\alpha A} \right)^{\frac{1}{t+1}}$$

and

$$\pi = \ln(1+r) + \frac{1}{1+r} \left(\frac{t+1}{\alpha A} \right)^{\frac{1}{t+1}} (\nu - \nu_c)^{\frac{1}{t+1}} \quad ,$$

which has a nontrivial power law behavior typical of second order critical phenomena.