

---

*Closed book and closed notes. No work needs to be shown.*

---

1. Which of the following statements is correct?
  - a. An object at rest could have a nonzero net force
  - b. An object in motion must have a net force on it
  - c. A nonzero net force on an object implies that it will move
  - d. A zero net force implies that an object will come to rest
  - e. None of the above
  
2. Block A is stacked on top of block B. What is the third law pair of the normal force on block A?
  - a. The force of gravity on block A
  - b. The normal force on block B
  - c. The friction force of block B on block A
  - d. The weight of block A on block B
  - e. There is no third law pair for this force.
  
3. Three forces act on an object with mass 1kg that is moving North with an acceleration of  $2.00 \text{ m/s}^2$ .  $\vec{F}_1 = 6.00 \text{ N}$  and it points West.  $\vec{F}_2 = 6.00 \text{ N}$  and it points South. What is the magnitude and direction of the third force?
  - a.  $\vec{F}_3 = 10.0 \text{ N}$  and it points  $53^\circ$  North of East
  - b.  $\vec{F}_3 = 10.0 \text{ N}$  and it points  $53^\circ$  South of West
  - c.  $\vec{F}_3 = 7.21 \text{ N}$  and it points  $34^\circ$  South of East
  - d.  $\vec{F}_3 = 7.21 \text{ N}$  and it points  $34^\circ$  North of East
  - e.  $\vec{F}_3 = 7.21 \text{ N}$  and it points  $56^\circ$  South of East
  
4. A football is kicked into the air at an angle  $\theta$  to the horizontal. What can be said about the net force on the football when it reaches its maximum height? Neglect air resistance.
  - a. The net force is zero.
  - b. The net force points horizontally.
  - c. The net force points upwards.
  - d. The net force is in the direction of the kick.
  - e. The net force points downwards.

5. You drag a 20.0 N box horizontally at a constant velocity with a rope that makes a  $30^\circ$  angle above the horizontal. What is the tension in the rope if the coefficient of kinetic friction between the box and the ground is 0.500?
- 16.2 N
  - 8.96 N
  - 11.5 N
  - 17.9 N
  - 40 N
6. Given that the mass of the moon is  $7.36 \times 10^{22}$  kg and the radius of the moon is 1737 km, how much would someone weigh on the moon if they weigh 600 N on Earth?
- 170 N
  - 1700 N
  - 600 N
  - 99.6 N
  - 976 N
7. A person who weighs 600 N on the ground is standing on a scale in an elevator that is descending with a velocity of 5 m/s and slowing down at  $2 \text{ m/s}^2$ . What is the reading on the scale?
- 600 N
  - 122 N
  - 722 N
  - 2400 N
  - 1200 N
8. A 5-kg concrete block is lowered with a downward acceleration of  $2.8 \text{ m/s}^2$  by means of a rope. The force of the block on the rope is:
- 14 N, up
  - 14 N, down
  - 35 N, up
  - 35 N, down
  - 49 N, up

9. A 10 kg block with an initial velocity of 10 m/s slides 10 meters across a horizontal surface and comes to rest. The magnitude of friction force of the surface acting on the block is:
- 5 N
  - 10 N
  - 25 N
  - 50 N
  - 98 N
10. A horizontal force of 12 N pushes a 0.50 kg book against a vertical wall. The book is initially at rest. If  $\mu_s = 0.60$  and  $\mu_k = 0.25$  which *one* of the following is true for this situation?
- The magnitude of the frictional force is 4.9 N.
  - The magnitude of the frictional force is 7.2 N.
  - The normal force of the wall on the book is 7.2 N.
  - The book will start moving downward and accelerate.
  - The book will start moving downward with a constant speed.
11. A horizontal applied force  $\vec{F}$  pushes on mass  $M$  which is placed next to a mass  $m$  such that they both move along a horizontal frictionless surface. The magnitude of the force of either of these blocks on the other is:
- $mF/(m + M)$
  - $mF/M$
  - $mF/(M - m)$
  - $MF/(M + m)$
  - $MF/m$
12. A car moves horizontally with a constant acceleration of  $3 \text{ m/s}^2$ . A ball is suspended by a string from the ceiling of the car. The ball does not swing, being at rest with respect to the car. What angle does the string make with the vertical.
- $17^\circ$
  - $35^\circ$
  - $52^\circ$
  - $73^\circ$
  - Cannot be found without knowing the length of the string

13. A 1000 kg car drives around a flat curve with a radius of 30 m. If the coefficient of static friction between the tires and the ground is  $\mu_s = 0.60$ , what is the maximum speed that the car can drive before the wheels start slipping? [*Approximate the car as a point mass, i.e. don't worry about the number of wheels it has or how it is shaped.*]
- a. 1.33 m/s
  - b. 1.76 m/s
  - c. 176 m/s
  - d. 420 m/s
  - e. 13.3 m/s
14. A mass  $m_1 = 20$  kg is on a frictionless incline and is connected by a rope and pulley to a mass  $m_2 = 10$  kg that is hanging vertically down the other side of the incline. At what angle of incline will the two masses be in equilibrium?
- a.  $30^\circ$
  - b.  $45^\circ$
  - c.  $0^\circ$
  - d.  $90^\circ$
  - e.  $60^\circ$
15. A 100 kg box is at a height of 5 m. It slides down a ramp that is at an incline of  $50^\circ$ . The coefficient of kinetic friction is 0.50. What is the final velocity of the box when it reaches the bottom of the ramp?
- a. 7.89 m/s
  - b. 6.29 m/s
  - c. 5.76 m/s
  - d. 7.53 m/s
  - e. 8.24 m/s

Equations and Constants

$$\left\{ \begin{array}{l} x = r \cos \theta \\ y = r \sin \theta \end{array} \right\}; \quad \left\{ \begin{array}{l} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1} \left( \frac{y}{x} \right) \end{array} \right\};$$

$$\left\{ \begin{array}{l} v_x = v_{0x} + a_x t \\ \Delta x = \frac{1}{2}(v_{0x} + v_x)t \\ \Delta x = v_{0x}t + \frac{1}{2}a_x t^2 \\ v_x^2 = v_{0x}^2 + 2a_x \Delta x \end{array} \right\}; \quad \left\{ \begin{array}{l} v_y = v_{0y} + a_y t \\ \Delta y = \frac{1}{2}(v_{0y} + v_y)t \\ \Delta y = v_{0y}t + \frac{1}{2}a_y t^2 \\ v_y^2 = v_{0y}^2 + 2a_y \Delta y \end{array} \right\};$$

$$\left\{ \begin{array}{l} \Delta x = x_f - x_i \\ \text{speed}_{\text{ave}} = \frac{d}{\Delta t} \end{array} \right\}; \quad \left\{ \begin{array}{l} v_{\text{ave}} = \frac{\Delta x}{\Delta t} \\ a_{\text{ave}} = \frac{\Delta v}{\Delta t} \end{array} \right\}; \quad \left\{ \begin{array}{l} v = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \\ a = \lim_{\Delta t \rightarrow 0} \frac{\Delta v}{\Delta t} \end{array} \right\};$$

$$\left\{ \begin{array}{l} a_c = \frac{v^2}{r} \\ T = \frac{2\pi r}{v} \end{array} \right\}; \quad x = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A};$$

$$\left\{ \begin{array}{l} \sum \vec{F} = 0 \rightarrow \vec{a} = 0 \\ \sum \vec{F} = m\vec{a} \\ \vec{F}_{2\text{on}1} = -\vec{F}_{1\text{on}2} \end{array} \right\}; \quad \left\{ \begin{array}{l} \sum F_x = ma_x \\ \sum F_y = ma_y \end{array} \right\}; \quad \left\{ \begin{array}{l} 0 \leq f_s \leq \mu_s F_N \\ f_k = \mu_k F_N \\ F_g = mg \\ F_d = bv \end{array} \right\};$$

$$\left\{ \begin{array}{l} F_g = G \frac{Mm}{r^2} \\ G = 6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2/\text{kg}^2 \\ g = 9.80 \text{ m/s}^2 \\ M_{\text{Earth}} = 5.98 \times 10^{24} \text{ kg} \\ R_{\text{Earth}} = 6.27 \times 10^6 \text{ m} \end{array} \right\};$$