

I.) Plasma on a Back-of-Envelope

→ Basic Ideas

→ What is a plasma?

- dilute gas
- charged particles - i.e. electrons + ions, usually protons - but net neutral
- long range Coulombic interaction
- screening

→ Recall - dilute gas

- basic ordering: (collisional transport)

$$d \ll r \ll l_{mfp} \ll L \quad \xrightarrow{\text{system size}}$$

range of force mean interparticle spacing mean free path
 (i.e. radius of hard sphere)

or long mean free path regime:

$$d \ll r \ll L \ll l_{mfp}$$

some key considerations:
estimates

$$\textcircled{1} \quad d^3 n = d^3 / r^3 \ll 1$$

\Rightarrow diluteness - particles \ominus free,
mostly non-interacting

\textcircled{2}

$$l_{\text{mfp}} \sim 1/n\tau \sim 1/n\pi d^2 \\ \sim \bar{r} (\bar{r}/d)^2$$

$$l_{\text{mfp}}/\bar{r} \sim (\bar{r}/d)^2$$

diluteness assures l_{mfp} exceeds
particle spacing (i.e. gas vs
liquid, etc.).

$$l_{\text{mfp}}/d \sim (\bar{r}/d)^3$$

diluteness assures l_{mfp} exceeds
range of force.

$$\textcircled{3} \quad v_c \sim v_{th} / l_{\text{mfp}} \sim v_{th} n \tau$$

\sim defines collision frequency.

For l_mfp :

Particles

∴

i.e. • CCCC

particle + interaction cylinder

$$V_{IC} \sim \pi L$$

or

$\alpha = \# \text{ collisions in cylinder of length } L$

$$\alpha = n \pi L$$

or

~~mean~~ mean length between collisions

$$\text{l}_\text{mfp} \approx L / \alpha = (\gamma n \pi)$$

or

$$\left(\frac{d\alpha}{dL} \right)^{-1} \approx 1 / n \pi \approx \text{l}_\text{mfp}$$

$\text{l}_\text{mfp} \sim \gamma n \pi$

(4) basic diffusivity:

$$D \sim v_{th} \ell m_p \sim v_{th}^2 / r_c$$

Now, for plasma:


- Force is Coulomb

$$\text{i.e. } V = e\phi \sim 1/r$$

\Rightarrow no scale associated!

\Rightarrow long range!

i.e. contract hard sphere



\therefore no d .

\Rightarrow screening occurs!

so now fundamental collisional scale ordering is:

$$\boxed{r < \lambda_D \ll l_{mfp} \ll L}$$

$r \sim \sqrt[3]{n}^{1/3} \rightarrow$ mean inter-particle spacing

$\lambda_D \sim \text{Debye length} \rightarrow$ key scale in plasma

$l_{mfp} \sim$ mean free path

$$l_{mfp} \sim \tau / n \sigma$$

Key: cross-section, with long range interaction


$L \rightarrow$ system size

key point: Debye length and?

For Debye Length:

in plasma

$$\frac{q}{4\pi}$$



test charge

plasma
adjusts
to screen

$$1/r \rightarrow e^{-r/\lambda_D}/r$$

charge
needs
energy
 $\rightarrow T$

$$\nabla^2 \phi = -4\pi \rho$$

$$= -4\pi n_0 e \left[\frac{\partial n_i}{\partial \phi} - \frac{\partial n_e}{\partial \phi} \right] + 4\pi \rho \frac{\partial (x-x)}{\partial \phi}$$

medium/plasma
response

$b = \infty$
 $h = \infty$

$$\partial n_i / n_0 = \exp \left[-k_B \phi / T_i \right]$$

$k_B \rightarrow 1$
eV

$$\partial n_e / n_0 = \exp \left[k_B \phi / T_e \right]$$

so, noting neutrality (net):

$$\nabla^2 \phi = -4\pi n_0 e \left[+ \frac{1 - \exp(-k_B \phi / T_i)}{T_i} - \frac{1}{T_e} + \frac{\exp(k_B \phi / T_e)}{T_e} \right]$$

$$\nabla^2 \phi = 4\pi n_0 e^2 \left(\frac{1}{T_e} + \frac{1}{T_i} \right) \phi$$

$$\phi = \exp[-r/\lambda_D] / r$$

$$1/\lambda_D^2 = 4\pi n_0 e^2 (1/T_e + 1/T_i)$$

Screening
Debye
Length

$$\lambda_D^2 = \left[4\pi n_0 e^2 (1/T_e + 1/T_i) \right]^{-1}$$

\Rightarrow

① Key features of plasma:

$$n \lambda_D^3 \gg 1$$

\Rightarrow large number of particles in Debye sphere

$$\lambda_D > r \sim 1/n^{1/3}$$

Why: \Rightarrow discreteness!

$$\Rightarrow T > e^2 / n$$

Z.

i.e. thermal energy must exceed electrostatic energy

$\Rightarrow \left\{ \begin{array}{l} \text{dilute} \\ \text{plasma, not crystal. !} \end{array} \right.$

check : $T > e^2/r$

$$\frac{nT}{ne^2} \bar{r} > 1$$

$$\Rightarrow \lambda_D^3 \bar{r} n > 1$$

$$\lambda_D^3 > \bar{r}^2 \quad \checkmark$$

N.B. : orders : $\bar{r} < \lambda_D$.

Also : plasma closed ;

why : Thermal Fluctuations exceed QM Fluctuations in energy.

Now : thermal $\rightarrow T$

$$-\nabla M \quad E \sim p^2/2m$$

$$\sim \hbar^2 r^2/2m$$

$$\sim \hbar^2/r^2 2m \sim \hbar^2 n^{2/3}/m$$

\Rightarrow

$$\boxed{T \gg \hbar^2 n^{2/3}/m}$$

and have:

$$T \gg e^2 n^{1/3} \quad \rightarrow \text{differences}$$

\approx , for dilute plasma:

$$e^2 n^{1/3} > \hbar^2 n^{2/3}/m$$

\Rightarrow

$$e^2 n^{1/3} / \hbar^2 n^{2/3}/m \sim \frac{me^2}{\hbar^2 n^{1/3}}$$

$$\simeq \frac{r}{a_B} \gg 1$$

mean interparticle
spacing must exceed
Bohr radius

$$a_B$$

Bohr radius

where: $a_B = \frac{4\pi\hbar^2}{me^2} \rightarrow$ Bohr radius

check: $\frac{\hbar^2}{2m a_B^2} \sim \frac{e^3}{a_B}$ $\sim 1 \text{ nm}$

$$a_B \sim \hbar^3/me^2$$

so conditions for $\rho \approx 0 \text{ m}^{-2}$:
(classical)

$$\boxed{n \lambda_0^3 \gg 1}$$

$$\lambda_0^3/a_B^3 \gg 1$$

and so have scale ordering:

$$\boxed{r < \lambda_0 < \text{lmp} < L}$$

Frequencies / Resonances

- much of plasma physics deals with waves / instabilities
- collective resonances

$$\underline{D} - \underline{D}_0 = 4\pi \rho_{ext} \quad \text{dielectric fn.}$$

$$\underline{D} = \underline{E} + 4\pi \underline{\rho} = G(\omega) \underline{E}$$

+ polarization

N.B. in most generality,
 $E(\omega) \rightarrow E(\omega, \mathbf{k})$

and, say, electron polarization:

$$\underline{\rho} = Ne \delta \underline{x} \quad (\text{ex})$$

i.e. consider high frequency wave oscillation, so electron inertia low.

$$m_e \frac{d^2 \underline{x}}{dt^2} = e \underline{E}$$

$$(E = E_0 e^{-i\omega t})$$

\Rightarrow \rightarrow excursion

$$-\omega_m^2 \underline{x} = e \underline{E}$$

$$\Rightarrow \underline{x} = -e \underline{E} / \omega_m^2 m_e$$

$$\stackrel{\text{so}}{=} 4\pi \underline{P} = -\frac{4\pi n_a e^2}{m_e \omega^2} \underline{E}$$

$$= -\omega_p^2 / \omega^2$$

$$\boxed{\omega_p^2 = \frac{4\pi n_a e^2}{m_e} \rightarrow \text{plasma frequency}}$$

\rightarrow space charge oscillation wave

$\rightarrow \delta n \rightarrow \delta E \Leftrightarrow$ restoring force.

$\stackrel{\text{so}}{=}$

$$\underline{D} = \underline{E} + 4\pi \underline{P} = \left(1 - \frac{\omega_p^2}{\omega^2}\right) \underline{E}$$

$$\epsilon(\omega) = 1 - \omega_p^2/\omega^2$$

50

$$\underline{D} \cdot \underline{D} = (1 - \omega_p^2/\omega^2) \underline{D} \cdot \underline{E} = 4\pi \rho_{ext}$$

50 → for $\rho_{ext} = \rho_{ext}$ ($\omega \sim \omega_p$)

⇒ E response is plasma
is large. $\Rightarrow E \rightarrow 0$

- collective resonance or mode


- origin is space charge separation.
⇒ restoring force.

identifies: $\omega_p^2 = 4\pi n e^2 / m$

fundamental of plasma frequency

Transport \leftrightarrow Coulomb Collisions

\rightarrow Consider Fermi/ker collision:



- what is cross section?

in particular seek cross section for weak deflection \leftrightarrow "momentum transfer cross-section"

↳ More glancing collisions occur...

- of course central force, so $|p|$ conserved but direction changed

$$m \Delta v_{\perp} = \Delta \Omega = \int_{-\infty}^{+\infty} dt \vec{F}_\perp$$

$$= \int_{-\infty}^{+\infty} dt \frac{e^2 \sin \alpha}{r^2}$$


$$= \int_{-\infty}^{+\infty} dt \frac{e^2}{r^2} \frac{\dot{\theta}}{r}$$

$$\Delta \theta_1 = e^2 \int_{-\infty}^{+\infty} \frac{dt}{(\rho^2 + v^2 t^2)^{3/2}} \sim e^2 \rho \int_{-\infty}^{+\infty} \frac{dt}{\rho^3 (1 + \frac{v^2 t^2}{\rho^2})^{3/2}}$$

$$\sim e^2 / \rho v$$

but $\Delta \theta_1 \sim u v \sin \theta$
 $\sim u v \theta$

so deflection angle:



$$\theta \sim e^2 / u v^2 \rho$$

then for cross section:

$$d\Gamma = \rho d\Omega = d(\rho^3) = d \left(\frac{e^2}{u v^2 \rho} \right)^2$$

↳ area of interaction cylinder

i.e. note key point: cross section
 heavily weights weak deflections

$$d\Gamma \approx \left(\frac{e^2}{u v^2} \right)^2 \frac{d\Omega}{\rho^3}$$

$$\text{Now: } d\Gamma = \left(\frac{e^2}{4\pi r^2}\right)^2 \frac{d\Omega}{r^3}$$

↑
weak deflection
divergence

A-h.: small $\theta \leftrightarrow$ large R

\Rightarrow long range character of Coulomb force!

\Rightarrow screening, long range cut-off is very relevant.

Neglect momentum transfer cross-section
need take out ~~collisions~~ collisions
with no transfer, i.e.

$$d\Gamma_f = (1 - \cos\theta) d\Gamma$$

$$\approx \theta^2 \left(\frac{e^2}{4\pi r^2}\right)^2 \frac{1}{r^3}$$

so $d\Gamma_f \approx \left(\frac{e^2}{4\pi r^2}\right)^2 \frac{1}{r^3}$

$$\tau_f \approx \left(\frac{e^2}{\mu v^2} \right)^2 \ln \left(\frac{1}{\theta_0} \right)$$

θ
divergence - low θ

- Coulomb cross-section, Rutherford.

- θ_0 is small angle cut-off

Now low $\theta \leftrightarrow$ large ρ

\Rightarrow small angle cut-off set by
large ρ

" " largest ρ can be no \leftrightarrow
screening limited!

Now,

$$\theta \sim e^2 / \mu v^2 \rho$$

θ_0

$$\theta_0 \sim e^2 / \mu v^2 \lambda_0$$

screening
cut-off

Q:

$$\text{So } \ln \frac{1}{\epsilon} = \ln \left(\frac{1}{e^2}\right) = \ln \left(\frac{T \alpha^2}{e^2}\right)$$

\downarrow

L (in LL) Coulomb

$$\boxed{\tau_+ \sim \left(\frac{e^2}{T}\right)^2 \ln \frac{1}{\epsilon}}$$

(can resolve by G \rightarrow L(B))
→ effective cross section

\Rightarrow

$$\boxed{\tau_+ \sim \bar{n}^2 \left(\frac{e^2}{\bar{n} T}\right)^2 \ln \frac{1}{\epsilon}}$$

Coulomb
cross
section

Note: $\left(\frac{e^2}{\bar{n} T}\right)^2 \rightarrow \left[\frac{1}{n \lambda_0^3}\right]^{2/3} \sim \frac{\bar{n}^{-4}}{\lambda_0^4}$

$$\left(\frac{\bar{n}^2}{\lambda_0^2}\right)^{-2} \sim \left(\frac{1}{n \lambda_0^3}\right)^{4/3}$$

$$\boxed{\tau_+ \sim \bar{n}^2 \left(\frac{1}{n \lambda_0^3}\right)^{4/3} \ln \frac{1}{\epsilon}}$$

Now,

$$l_{\text{mfp}} \sim \frac{1}{n \tau_f}$$

$$\sim \frac{1}{n r^2} \left(\frac{e^2}{\pi T} \right)^2 \ln 1$$

$$\sim r \left(n \lambda_0^3 \right)^{4/3} / \ln 1$$

$$l_{\text{mfp}} \sim r \left(\lambda_0 / r \right)^4 / \ln 1$$

so $l_{\text{mfp}} \sim r \left(\lambda_0 / r \right)^4 / \ln 1$

$$\underline{l_{\text{mfp}}} \sim \left(\frac{\lambda_0}{r} \right)^3 / \ln 1$$

$$\sim n \lambda_0^3 / \ln 1$$

as $n \lambda_0^3 \gg \ln 1$

$$l_{\text{mfp}} \gg \lambda_0$$

consistent
with screening

so

$$r < \lambda_D < l_{mfp} < L$$

collisional plasma ordering

Note:

$$\frac{l_{mfp}}{\lambda_D} \approx \frac{v}{\lambda_D} \left(\frac{\lambda_D}{r} \right)^4 / \ln \Lambda$$

$$\approx (n \lambda_D^3) / \ln \Lambda$$

Now, further points about transport:

- apart from $\ln \Lambda$, no mass, n scaling in T_f , l_{mfp} .

- so $\tau_{col} \sim M^{-1/2} \rightarrow$ via V_{th} .

$$\tau_{col} \sim M^{-1/2} \quad \text{via } V_{th}.$$

$$\tau_{ee} \sim (M_e/M_i)^{1/2} \ll 1.$$

20
1.

then, as before (gives) :

→ thermal conductivity

$$\lambda \sim n v_{th} \text{ lmax}$$

\downarrow } \downarrow
 C_V } index α

longer for electrons

so

electrons control thermal conduction

→ viscosity

$$\eta \sim M_i n v_{th} \text{ lmax}$$

control flow

$$\text{lmax} \sim 1/n\tau$$

contrast:

$$\eta \sim M_i n v_{th} \text{ lmax}$$

$$\boxed{\eta \sim M_i V_{thc} / T_F}$$

$$\boxed{\eta \sim (M_i T)^{1/2} / T_F}$$