

$$23. \quad (a) \quad E_0 = \frac{1}{2} \hbar \omega_0 = \frac{1}{2} k x_0^2 \quad \text{so} \quad x_0 = \sqrt{\hbar \omega_0 / k}$$

$$(b) \quad E_1 = \frac{3}{2} \hbar \omega_0 = \frac{1}{2} k x_0^2 \quad \text{so} \quad x_0 = \sqrt{3 \hbar \omega_0 / k}$$

$$E_2 = \frac{5}{2} \hbar \omega_0 = \frac{1}{2} k x_0^2 \quad \text{so} \quad x_0 = \sqrt{5 \hbar \omega_0 / k}$$

$$24. \quad x_{\text{av}} = \int_{-\infty}^{\infty} |\psi(x)|^2 x dx = A^2 \int_{-\infty}^{\infty} e^{-2ax^2} x dx = 0$$

because the integrand is an odd function of  $x$  (the integral from  $-\infty$  to  $0$  exactly cancels the integral from  $0$  to  $+\infty$ ).

$$(x^2)_{\text{av}} = \int_{-\infty}^{\infty} |\psi(x)|^2 x^2 dx = A^2 \int_{-\infty}^{\infty} e^{-2ax^2} x^2 dx = 2A^2 \int_0^{\infty} e^{-2ax^2} x^2 dx = \frac{2A^2}{\sqrt{8a^3}} \int_0^{\infty} e^{-u^2} u^2 du$$

with the substitution  $u = x\sqrt{2a}$ . The integral is a standard form found in tables and is equal to  $\sqrt{\pi}/4$ . Substituting  $A = (\omega_0 m / \pi \hbar)^{1/4}$  and  $a = \sqrt{km}/2\hbar = \omega_0 m / 2\hbar$ , we find

$$(x^2)_{\text{av}} = 2 \left( \frac{\omega_0 m}{\pi \hbar} \right)^{1/2} \frac{1}{2\sqrt{2}} \left( \frac{2\hbar}{\omega_0 m} \right)^{3/2} \frac{\sqrt{\pi}}{4} = \frac{\hbar}{2\omega_0 m}$$

$$\Delta x = \sqrt{(x^2)_{\text{av}} - (x_{\text{av}})^2} = \sqrt{\hbar / 2m\omega_0}$$

25. (a) Because the oscillating particle moves with equal probability in the positive and negative  $x$  directions,  $p_{\text{av}} = 0$ .

$$(b) \quad U_{\text{av}} = \frac{1}{2} k (x^2)_{\text{av}} = \frac{1}{2} k \frac{\hbar}{2\omega_0 m} = \frac{1}{2} \omega_0^2 m \frac{\hbar}{2\omega_0^2 m} = \frac{1}{4} \hbar \omega_0$$

$$K_{\text{av}} = E - U_{\text{av}} = \frac{1}{2} \hbar \omega_0 - \frac{1}{4} \hbar \omega_0 = \frac{1}{4} \hbar \omega_0$$

$$(p^2)_{\text{av}} = 2mK_{\text{av}} = 2m \left( \frac{1}{4} \hbar \omega_0 \right) = \frac{\hbar \omega_0 m}{2}$$

$$(c) \quad \Delta p = \sqrt{(p^2)_{\text{av}} - (p_{\text{av}})^2} = \sqrt{\hbar\omega_0 m/2}$$

$$26. \quad E_0 = 1.24 \text{ eV} = \frac{1}{2}\hbar\omega_0 \quad \text{so} \quad \hbar\omega_0 = 2.48 \text{ eV}$$

$$\text{To } n = 2 \text{ state:} \quad \Delta E = E_2 - E_0 = \frac{5}{2}\hbar\omega_0 - \frac{1}{2}\hbar\omega_0 = 2\hbar\omega_0 = 2(2.48 \text{ eV}) = 4.96 \text{ eV}$$

$$\text{To } n = 4 \text{ state:} \quad \Delta E = E_4 - E_0 = \frac{9}{2}\hbar\omega_0 - \frac{1}{2}\hbar\omega_0 = 4\hbar\omega_0 = 4(2.48 \text{ eV}) = 9.92 \text{ eV}$$

$$27. \quad P(x) dx = |\psi(x)|^2 dx = A^2 e^{-2ax^2} dx \quad \text{so at } x = 0 \quad P(0)dx = A^2 dx$$

At the classical turning points  $x = \pm x_0$ ,  $K = 0$  so  $E = U$  or  $\frac{1}{2}\hbar\omega_0 = \frac{1}{2}kx_0^2$

$$P(\pm x_0)dx = A^2 e^{-2(\sqrt{km}/2\hbar)(\hbar\omega_0/k)} dx = A^2 e^{-1} dx = e^{-1} P(0)dx = 0.368 P(0)dx$$

28. (a) If  $E = 0$ , then  $p = 0$  and we would know the momentum exactly. Thus  $\Delta p = 0$ , which means  $\Delta x = \infty$ . But that would be inconsistent with a particle that is bound to a finite region of space.

(b)

$$E = \frac{1}{2}\hbar\omega_0 = \frac{1}{2}\hbar\sqrt{\frac{k}{m}} = \frac{1}{2}\hbar c\sqrt{\frac{k}{mc^2}} = 0.5(197 \text{ eV} \cdot \text{nm})\sqrt{\frac{3.5 \times 10^3 \text{ eV/nm}^2}{938 \times 10^3 \text{ eV}}} = 0.19 \text{ eV}$$

This is less than the binding energy, so this motion is not sufficient to dissociate the molecule.

(c) At the turning point of the motion,  $E = \frac{1}{2}kx_0^2$ , so

$$x_0 = \sqrt{\frac{2E}{k}} = \sqrt{\frac{2(0.19 \text{ eV})}{3.5 \times 10^3 \text{ eV/nm}^2}} = 0.010 \text{ nm}$$

This motion is not negligible at the atomic level.

$$32. \quad x < 0: \quad \psi_0 = A'e^{ik_0x} + B'e^{-ik_0x} \quad \text{with} \quad k_0 = \sqrt{\frac{2mE}{\hbar^2}}$$

$$x > 0: \quad \psi_1(x) = C'e^{ik_1x} + D'e^{-ik_1x} \quad \text{with} \quad k_1 = \sqrt{\frac{2m(E - U_0)}{\hbar^2}}$$

If the particles are incident from the negative  $x$  direction, then  $D'$  (which is the coefficient of the term that represents a wave in the region of positive  $x$  traveling toward the origin) must be set to 0. We then apply the continuity conditions on  $\psi$  and  $d\psi/dx$  at  $x = 0$ :

$$\psi_0(0) = \psi_1(0): \quad A' + B' = C'$$

$$\left(\frac{d\psi_0}{dx}\right)_{x=0} = \left(\frac{d\psi_1}{dx}\right)_{x=0} : \quad k_0(A' - B') = k_1C'$$

39. (a) The particle has no preferred direction of motion, so it is equally likely to be moving in the positive and negative  $x$  directions. We therefore expect that  $p_{\text{av}} = 0$ .
- (b) Because the potential energy is zero inside the well, the kinetic energy is equal to the total energy:

$$K = E_n \quad \text{or} \quad \frac{p^2}{2m} = \frac{h^2 n^2}{8mL^2} \quad \text{so} \quad p^2 = \frac{h^2 n^2}{4L^2}$$

For a given level  $n$ ,  $p^2$  is constant so  $(p^2)_{\text{av}}$  has that same value.

$$(c) \quad \Delta p = \sqrt{(p^2)_{\text{av}} - (p_{\text{av}})^2} = \sqrt{\frac{h^2 n^2}{4L^2} - 0} = \frac{hn}{2L}$$

42. (a) The  $x$  and  $y$  motions are independent, and each contributes an energy of  $\hbar\omega_0(n + \frac{1}{2})$ , but the integer  $n$  is not necessarily the same for the two independent motions. Thus the total energy is

$$E = \hbar\omega_0(n_x + \frac{1}{2}) + \hbar\omega_0(n_y + \frac{1}{2}) = \hbar\omega_0(n_x + n_y + 1)$$

(b)

$4\hbar\omega_0$		4	(0,3), (1,2), (2,1), (3,0)
$3\hbar\omega_0$		3	(0,2), (1,1), (2,0)
$2\hbar\omega_0$		2	(0,1), (1,0)
$\hbar\omega_0$		1	(0,0)
Energy		Degeneracy	( $n_x, n_y$ )

(c) The level with energy  $N\hbar\omega_0$  has  $N$  different possible sets of quantum numbers  $n_x, n_y$ . Both  $n_x$  and  $n_y$  range from 0 to  $N-1$  but with their sum fixed to  $N$ . The number of possible values of  $n_x$  is then  $N$  (the values are 0, 1, 2, ...,  $N-2$ ,  $N-1$ ), and for each value of  $n_x$  the value of  $n_y$  is fixed. The total degeneracy of each level is thus  $N = n_x + n_y + 1$ .

43. (a) With  $\Delta x = \sqrt{(x^2)_{\text{av}} - (x_{\text{av}})^2}$ , clearly  $x_{\text{av}} = 0$  for this wave function. Then

$$(x^2)_{\text{av}} = \int_{-\infty}^{+\infty} x^2 |\psi(x)|^2 dx = 2b^{-1} \int_0^{+\infty} x^2 e^{-2x/b} dx = 2b^{-1} \frac{2}{(2/b)^3} = \frac{b^2}{2}$$

So  $\Delta x = b/\sqrt{2} = 0.71b$ .

(b) The maximum probability density occurs at  $x = 0$ , where  $P(x) = |\psi(x)|^2 = b^{-1}$ . We now find the location where  $P(x)$  drops to half that value, that is, where  $e^{-2|x|/b} = 0.5$ , or  $-2|x|/b = \ln(0.5)$ :

$$|x| = -(b/2)\ln(0.5) \quad \text{or} \quad x = \pm 0.347b$$

Our estimate for  $\Delta x$  is then the distance between the two points where the probability is half its maximum value, so  $\Delta x = 0.69b$ , which agrees very well with the result of the more rigorous calculation from part (a).