

**Justify all your answers to all problems. Write clearly.**

Time dilation; Length contraction:  $\Delta t = \gamma \Delta t_0$  ;  $L = L_0 / \gamma$  ;  $c = 3 \times 10^8 \text{ m/s}$

Lorentz transformation:  $x' = \gamma(x - ut)$  ;  $y' = y$  ;  $t' = \gamma(t - ux/c^2)$

Velocity:  $v'_x = \frac{v_x - u}{1 - uv_x/c^2}$  ;  $v'_y = \frac{v_y}{\gamma(1 - uv_x/c^2)}$  ;  $\gamma = \frac{1}{\sqrt{1 - u^2/c^2}}$

Inverse transformations:  $u \rightarrow -u$ , primed  $\leftrightarrow$  unprimed; Doppler:  $f' = f \sqrt{\frac{1 \pm u/c}{1 \mp u/c}}$

Momentum:  $\vec{p} = \gamma m \vec{v}$  ; Energy:  $E = \gamma mc^2$  ; Kinetic energy:  $K = (\gamma - 1)mc^2$   
 $E = \sqrt{p^2 c^2 + m^2 c^4}$  ; rest energy:  $E_0 = mc^2$

Electron:  $m_e = 0.511 \text{ MeV}/c^2$  ; Proton:  $m_p = 938.26 \text{ MeV}/c^2$  ; Neutron:  $m_n = 939.55 \text{ MeV}/c^2$

Atomic unit:  $1u = 931.5 \text{ MeV}/c^2$  ; electron volt:  $1\text{eV} = 1.6 \times 10^{-19} \text{ J}$

Photoelectric effect:  $eV_s = K_{\text{max}} = hf - \phi = hc/\lambda - \phi$  ;  $\phi$  = work function

Stefan law:  $I = \sigma T^4$  ,  $\sigma = 5.67037 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$  ; Wien's law:  $\lambda_m T = 2.8978 \times 10^{-3} \text{ m} \cdot \text{K}$

$I(T) = \int_0^\infty I(\lambda, T) d\lambda$  ;  $I = (c/4)u$  ;  $u(\lambda, T) = N(\lambda)E_{\text{av}}(\lambda, T)$  ;  $N(\lambda) = \frac{8\pi}{\lambda^4}$

Boltzmann distribution:  $N(E) = Ce^{-E/kT}$  ;  $N = \int_0^\infty N(E) dE$  ;  $E_{\text{av}} = \frac{1}{N} \int_0^\infty EN(E) dE$

Classical:  $E_{\text{av}} = kT$  ; Planck:  $E_n = n\varepsilon = nhf$  ;  $N = \sum_{n=0}^\infty N(E_n)$  ;  $E_{\text{av}} = \frac{1}{N} \sum_{n=0}^\infty E_n N(E_n)$

Planck:  $E_{\text{av}} = \frac{hc/\lambda}{e^{hc/\lambda kT} - 1}$  ;  $hc = 1,240 \text{ eV} \cdot \text{nm}$  ;  $\lambda_m T = hc/4.96k$  ;  $\sigma = \frac{2\pi^5 k^4}{15c^2 h^3}$

Boltzmann constant:  $k = (1/11,604) \text{ eV}/\text{K}$  ;  $1\text{\AA} = 1\text{A} = 0.1 \text{ nm}$

Compton scattering:  $\lambda' - \lambda = \lambda_c(1 - \cos\theta)$  ;  $\lambda_c = \frac{h}{m_e c} = 0.00243 \text{ nm}$

double-slit interference maxima:  $d \sin\theta = n\lambda$  ; single-slit diffraction minima:  $a \sin\theta = n\lambda$

de Broglie:  $\lambda = \frac{h}{p}$  ;  $f = \frac{E}{h}$  ;  $\omega = 2\pi f$  ;  $k = \frac{2\pi}{\lambda}$  ;  $E = \hbar\omega$  ;  $p = \hbar k$  ;  $\hbar c = 197.3 \text{ eV nm}$

matter:  $E = \frac{p^2}{2m}$  (nonrelativistic) or  $E = \sqrt{p^2 c^2 + m^2 c^4}$  (relativistic); photons:  $E = pc$

Uncertainty:  $\Delta x \Delta k \sim 1$  ;  $\Delta t \Delta \omega \sim 1$  ;  $\Delta x \Delta p \sim \hbar$  ;  $\Delta t \Delta E \sim \hbar$  ;  $\Delta A = \sqrt{\langle A^2 \rangle - \langle A \rangle^2}$

group and phase velocity :  $v_g = \frac{d\omega}{dk}$  ;  $v_p = \frac{\omega}{k}$  ; wave packets:  $\psi(x,t) = \int a(k) e^{i(kx - \omega(k)t)}$

Schrodinger equation :  $-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + U(x)\Psi(x,t) = i\hbar \frac{\partial \Psi}{\partial t}$  ;  $\Psi(x,t) = \psi(x)e^{-i\frac{E}{\hbar}t}$

Time - independent Schrodinger equation :  $-\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + U(x)\psi(x) = E\psi(x)$  ;  $\int_{-\infty}^{\infty} dx \psi^* \psi = 1$

$P(x_1 < x < x_2) = \int_{x_1}^{x_2} |\psi(x)|^2 dx$  ;  $\langle f(x) \rangle = \int_{-\infty}^{\infty} |\psi(x)|^2 f(x) dx$

$\infty$  square well:  $\psi_n(x) = \sqrt{\frac{2}{L}} \sin(\frac{n\pi x}{L})$  ;  $E_n = \frac{\pi^2 \hbar^2 n^2}{2mL^2}$  ;  $\frac{\hbar^2}{2m_e} = 0.0381 \text{ eV nm}^2$  (electron)

2D square well:  $\Psi_{n_1 n_2}(x,y) = \Psi_{1,n_1}(x)\Psi_{2,n_2}(y)$  ;  $E_{n_1 n_2} = \frac{\pi^2 \hbar^2}{2m} (\frac{n_1^2}{L_1^2} + \frac{n_2^2}{L_2^2})$  ;  $\Psi_{i,n}(w) = \sqrt{\frac{2}{L_i}} \sin(\frac{n\pi w}{L_i})$

Harmonic oscillator:  $\Psi_n(x) = H_n(x)e^{-\frac{m\omega}{2\hbar}x^2}$  ;  $E_n = (n + \frac{1}{2})\hbar\omega$  ;  $E = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2 = \frac{1}{2}m\omega^2 A^2$

Step potential: reflection coef :  $R = \frac{(k_1 - k_2)^2}{(k_1 + k_2)^2}$  ,  $T = 1 - R$  ;  $k = \sqrt{\frac{2m}{\hbar^2}(E - U)}$

Tunneling:  $\psi(x) \sim e^{-\alpha x}$  ;  $T = e^{-2\alpha \Delta x}$  ;  $T = e^{-2 \int_{x_1}^{x_2} \alpha(x) dx}$  ;  $\alpha(x) = \sqrt{\frac{2m[U(x) - E]}{\hbar^2}}$

Rutherford scattering:  $U = \frac{2Ze^2}{4\pi\epsilon_0 r}$  ;  $e^2/(4\pi\epsilon_0) = 1.44 \text{ eV} \cdot \text{nm}$

$b = \frac{Z}{K_\alpha} \frac{e^2}{4\pi\epsilon_0} \cot \frac{1}{2}\theta$  ;  $f_{>\theta} = n\pi b^2$  ;  $N(\theta) = \text{constant} \times (\frac{Z}{K_\alpha})^2 \times \frac{1}{\sin^4(\theta/2)}$

Line spectra:  $\frac{1}{\lambda} = R(\frac{1}{n_0^2} - \frac{1}{n^2})$  ;  $R = \frac{1}{91.13 \text{ nm}}$  ;  $hcR = 13.6 \text{ eV}$

Bohr atom:  $F = \frac{ke^2 Z}{r^2} = m_e \frac{v^2}{r}$  ;  $U = -\frac{ke^2 Z}{r}$  ;  $E = K + U = -\frac{ke^2 Z}{2r}$  ;  $a_0 = 0.0529 \text{ nm}$

$r_n = (a_0 / Z)n^2$  ;  $E_n = -E_0 Z^2 / n^2$  ;  $E_0 = \frac{ke^2}{2a_0}$  ;  $a_0 = \frac{\hbar^2}{m_e ke^2}$  ;  $L = m_e v r = n\hbar$  ;  $E_0 = -13.6 \text{ eV}$

1-dim atom:  $U(x) = -ke^2 / x$  ; ground state wavefunction  $\psi(x) = Ax e^{-x/a_0}$

Spherically symmetric potential:  $\Psi_{n,\ell,m_\ell}(r,\theta,\phi) = R_{n\ell}(r)Y_{\ell,m_\ell}(\theta,\phi)$  ;  $Y_{\ell,m_\ell}(\theta,\phi) = P_\ell^{m_\ell}(\theta)e^{im_\ell\phi}$

quantum numbers:  $n = 1, 2, 3, \dots$  ;  $0 \leq \ell \leq n - 1$  ;  $-\ell \leq m_\ell \leq \ell$

Angular momentum:  $\vec{L} = \vec{r} \times \vec{p}$  ;  $|L^2| = \ell(\ell+1)\hbar^2$  ;  $L_z = m_\ell \hbar$

Radial probability density:  $P(r) = r^2 |R_{n\ell}(r)|^2$  ; Energy:  $E_n = -\frac{ke^2}{2a_0} \frac{Z^2}{n^2}$

Radial wave function:  $R_{n\ell}(r) = (\text{n-th degree polynomial in } r) \times e^{-Zr/na_0}$

Ground state of hydrogen-like ions:  $\Psi_{1,0,0} = \frac{1}{\pi^{1/2}} \left(\frac{Z}{a_0}\right)^{3/2} e^{-Zr/a_0}$  ;  $\int_0^\infty dr r^n e^{-\lambda r} = \frac{n!}{\lambda^{n+1}}$

Orbital magnetic moment:  $\vec{\mu} = \frac{-e}{2m_e} \vec{L}$  ;  $\mu_z = -\mu_B m_\ell$  ;  $\mu_B = \frac{e\hbar}{2m_e} = 5.79 \times 10^{-5} \text{ eV/T}$

Spin 1/2:  $s = \frac{1}{2}$  ,  $|\vec{S}| = \sqrt{s(s+1)}\hbar$  ;  $S_z = m_s \hbar$  ;  $m_s = \pm 1/2$  ;  $\vec{\mu}_s = \frac{-e}{2m_e} g\vec{S}$  ;  $g = 2$

Orbital+spin mag moment:  $\vec{\mu} = \frac{-e}{2m_e} (\vec{L} + g\vec{S})$  ; Energy in mag. field:  $U = -\vec{\mu} \cdot \vec{B}$

subshells: 1s, 2s, 2p, 3s, 3p, 4s, 3d, 4p, 5s, 4d, 5p, 6s, 4f, 5d, 6p, 7s, 5f, 6d

$2(2\ell + 1)$  electrons per subshell. Screened electron:  $E_n = (-13.6 \text{ eV}) Z_{\text{eff}}^2 / n^2$